

A UNIFIED APPROACH TO COMPACT SYMMETRIC SPACES OF RANK ONE

ADAM KORÁNYI AND FULVIO RICCI

ABSTRACT. A relatively simple algebraic framework is given, in which all the compact symmetric spaces can be described and handled without distinguishing cases. We also give some applications and further results.

INTRODUCTION

This article deals with compact Riemannian globally symmetric spaces of rank one. These spaces are very important for geometry, and of course they are well known. They are the spheres and the projective spaces over $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ (in the last case only of dimension ≤ 2).

There are two standard ways of describing them and make computations on them. One is to use classification; this is the approach, e.g., in [Be]. The projective spaces over $\mathbb{R}, \mathbb{C}, \mathbb{H}$ are easily handled together, but the case of \mathbb{O} is different because of the non-associativity of the octonions. For the \mathbb{O} -case one has to refer, as does [Be], to the rather complicated articles of Freudenthal (cf., e.g., [Ba, S]).

The other way is to use the general theory of symmetric spaces [Hel1]. Here there is no need to distinguish cases, but one has to use the large machinery of semisimple Lie group theory, which was designed for much more general situations and is rather unwieldy when applied to the special case of spaces of rank one.

In this article we set up a new framework which does not make explicit use of the octonions and applies to spaces of rank one in an essentially unified way and makes it easy to work on them, including the \mathbb{O} -case. This is done in the first six sections.

The basic notion is that of a *C-module with J^2 -condition* (briefly: J^2C -module). A *C-module* is the same thing as the classical notion of composition of quadratic forms [Hus], and is closely related to the notion of a Clifford module. The J^2 -condition specifies a subclass; it already played a fundamental rôle in [CDKR2]. As we have discovered since, its basic idea goes back to [Hei].

For non-compact symmetric spaces, a program similar to the present one was carried out in [CDKR2] and our article can be regarded as a continuation of [CDKR2], although our starting point is somewhat different. In [CDKR2] the basic objects are *H-type Lie algebras*; they form a category essentially equivalent to that of Clifford modules.

Research partially supported by the National Science Foundation, a PSC-CUNY grant, Centro de Giorgi, Pisa and GNAMPA-INdAM, Rome.

Some of the arguments in the introductory sections of this article are classical or are reformulations of proofs in [CDKR2], but we prove everything we need. Some other facts, which we do not need but which are of interest for understanding the full picture, are only described without proof in Section 3. This section also contains the classification of J^2C -modules, with proofs. This, together with the known classification of all symmetric spaces [Hel1, W], could be used to quickly show that our theory covers all the compact rank-one symmetric spaces. However, we will give a classification-free proof of this fact in Section 8.

Sections 1 and 2 contain the basic properties of J^2C -modules. In Section 4 we introduce the space $W = C \oplus V$, with V a J^2C -module. W is a weaker substitute for a J^2C -module, but a good notion of C -line can still be defined in it.

With these tools, we construct CPW as a compactification of W by adjoining a point at infinity for every family of parallel C -lines in W . In Section 5 we describe the topology and the differentiable structure on CPW , and in Section 6 its metric and its isometry group U , proving that CPW actually is a compact rank-one symmetric space. This ends the construction of the symmetric spaces.

In the subsequent part of the paper we prove various facts about the geometry of these spaces and the relevant transformation groups on them.

In Section 7 we illustrate the usefulness of our construction by reproving in simple ways some known properties of these spaces. As mentioned already, we prove in Section 8 that they are all the compact rank-one symmetric spaces, without appealing to classification.

In Section 9 we analyze the structure of the group $GL(W, C)$ of invertible linear transformations of W which preserve C -lines. This is preliminary to Section 10, where we introduce the group \mathcal{G} of collineations of CPW . \mathcal{G} is a transformation group properly containing the isometry group U and $GL(W, C)$, and whose elements are characterized by the property of preserving projective C -lines (i.e. closures in CPW of the C -lines of W , or images of these under isometries). We prove that collineations form a semisimple Lie group and that they act conformally on each projective C -line.

In Section 11 we prove that the collineation groups \mathcal{G} are characterized as the semisimple parts of automorphism groups of irreducible symmetric cones. We do so by constructing a representation of \mathcal{G} and identifying the cone in a real form of the representation space. The induced projective action of \mathcal{G} on the space of extremal lines of the cone provides an identification of this space with CPW .

In the Appendix at the end of the paper we show how the general non-compact symmetric space can be realized as the unit ball in W with a different metric. This amounts to redoing [CDKR2] from a different starting point.

We wish to point out that our construction results in an actual unification in the geometric as well as the algebraic sense. The spheres are included as the extreme case $V = 0$ and C arbitrary. In such a case the whole CPW consists of a single projective C -line. The other extreme case, $C = \mathbb{R}$ and V arbitrary, gives real projective space. In both extreme cases the unit ball of W is real hyperbolic space; in the first case it appears as the Poincaré model, in the second as the Klein model. In the cases in between, it agrees with the models used in [M].

TABLE OF CONTENTS

1. C -modules with the J^2 -condition

2. Automorphisms of J^2C -modules
3. Classification and some background information
4. The space $W = C \oplus V$
5. The construction of CPW
6. CPW as a compact symmetric space
7. Some applications
8. Every compact rank-one space is a CPW
9. The group $GL(W, C)$
10. The group of collineations
11. Compact rank-one spaces and symmetric cones
- A. Appendix. The non-compact symmetric spaces

1. C -MODULES WITH THE J^2 -CONDITION

Let C, V be finite-dimensional real Euclidean spaces over \mathbb{R} . We write the inner products as $\langle \cdot, \cdot \rangle$ and the norms as $|\cdot|$. We assume that C has a distinguished unit element, denoted by $\mathbf{1}$.

A C -module structure on V is a bilinear map $J : C \times V \rightarrow V$, such that

$$(1.1) \quad J(\mathbf{1}, v) = v \quad (\forall v \in V),$$

$$(1.2) \quad |J(\zeta, v)| = |\zeta||v| \quad (\forall \zeta \in C, v \in V).$$

Instead of $J(\zeta, v)$ we will also use the notation $J_\zeta v$ and, more frequently, ζv . We also write $Cv = \{\zeta v : \zeta \in C\}$ for $v \in V$.

We say that the C -module V satisfies the J^2 -condition (or, briefly, is a J^2C -module) if

$$(1.3) \quad C(Cv) = Cv \quad (\forall v \in V).$$

We do not exclude the “trivial” cases where $C = \mathbb{R}\mathbf{1}$ or $V = 0$. Notice that in such cases the J^2 -condition is trivially satisfied and most of the next definitions are vacuous.

We denote by C' the orthogonal complement of $\mathbb{R}\mathbf{1}$ in C . If $\zeta = a\mathbf{1} + z$, with $a \in \mathbb{R}$ and $z \in C'$, we set $\bar{\zeta} = a\mathbf{1} - z$, $a = \Re \zeta$, $z = \Im \zeta$.

Polarizing (1.2) in both ζ and v , we have

$$(1.4) \quad \langle \zeta u, \eta v \rangle + \langle \eta u, \zeta v \rangle = 2\langle \zeta, \eta \rangle \langle u, v \rangle,$$

Taking $\eta = \mathbf{1}$, $\zeta = z \in C'$, we see that J_z is skew-symmetric. For general $\zeta \in C$, this implies that

$$(1.5) \quad J_{\bar{\zeta}} J_\zeta = |\zeta|^2 \text{id}.$$

For $z \in C'$, (1.5) gives $J_z^2 = -|z|^2 \text{id}$. From this it is clear that the action of C' on V via J extends to an action of the Clifford algebra $\text{Cliff}(C')$. (We recall that $\text{Cliff}(C')$ is the associative algebra generated by C' and a unit element e subject to the relations $z^2 = -|z|^2 e$. In other words, if $\{z_1, \dots, z_m\}$ is an orthonormal basis of C' , then $\text{Cliff}(C')$ is generated by these elements and e , subject to the relations $z_i z_j + z_j z_i = -2\delta_{ij} e$.)

So V is automatically a *Clifford module*, meaning a representation of the associative algebra $\text{Cliff}(C')$. (More exactly we should say an “orthogonal” Clifford module, to take into account the added condition $|zv| = |z||v|$ for all $z \in C'$, $v \in V$.)

For any $\alpha \in \text{Cliff}(C')$, we still write its action on $v \in V$ as αv .

By (1.3), for every $v \neq 0$ in a J^2C -module V , Cv is a $\text{Cliff}(C')$ submodule, necessarily irreducible. From (1.4), it follows that the orthogonal complement of a submodule is again a submodule. So we can inductively construct *orthonormal C -bases*, i.e. orthonormal sets $\{v_1, \dots, v_n\}$ such that V is the orthogonal sum of Cv_1, \dots, Cv_n . If $V = Cv$ for some $v \neq 0$, we say that the J^2C -module is irreducible.

Another important property of J^2C -modules is that any non-zero element $v \in V$ determines a multiplication law on C , denoted by \cdot_v and given by

$$(\zeta \cdot_v \eta)v = \zeta\eta v .$$

Obviously, $\cdot_{tv} = \cdot_v$ for $t \in \mathbb{R}^*$.

For the remainder of this section, we assume that V is a non-trivial J^2C -module (but we allow $C = \mathbb{R}\mathbf{1}$).

Proposition 1.1. *Under the multiplication \cdot_v , C is a normed division algebra.*

Proof. The product is well defined because the map $\zeta \mapsto \zeta v$ from C to Cv is a bijection. It is obvious that the product is bilinear and has $\mathbf{1}$ as its identity element. Also, $|\zeta \cdot_v \eta| = |\zeta||\eta|$ by (1.2). To show that C is a division algebra, we must solve the equations

$$\xi \cdot_v \eta = \zeta , \quad \eta \cdot_v \xi' = \zeta$$

in ξ, ξ' , when ζ and $\eta \neq 0$ are given. For the first equation, we have to solve

$$\xi\eta v = \zeta v .$$

This can be done, because the map $\lambda \mapsto \lambda(\eta v)$ is a bijection of C onto $C(\eta v) \subset C(Cv) = Cv$. Since both Cv and $C(\eta v)$ have the same dimension as C , we have $C(\eta v) = Cv$.

To solve the equation

$$\eta\xi'v = \zeta v ,$$

we multiply both sides by $\bar{\eta}$ on the left and use (1.5) to obtain that

$$\xi'v = |\eta|^{-2}\bar{\eta}\zeta v ,$$

i.e. $\xi' = |\eta|^{-2}\bar{\eta} \cdot_v \zeta$. \square

Remark. As we shall see next in detail, \cdot_v can actually depend on v , and this dependence is equivalent to lack of associativity. Nevertheless, some expressions are independent of v in general. We list a few.

1. The proof of Proposition 1.1 shows that $\zeta^{-1} = |\zeta|^{-2}\bar{\zeta}$ when $\zeta \neq 0$, with respect to any multiplication \cdot_v . It follows that the value of any other rational expression in ζ with real coefficients is independent of v and coincides with its value with respect to Clifford multiplication.

2. For any $\zeta, \eta \in C$, the product $\zeta \cdot_v (\eta \cdot_v \zeta)$ equals the Clifford algebra product $\zeta \eta \zeta$ and therefore it does not depend on v . To see this, it is convenient to consider separately the case $\eta \in \mathbb{R}\mathbf{1} + \mathbb{R}\Im \zeta$ and the case $\eta \perp (\mathbb{R}\mathbf{1} + \mathbb{R}\Im \zeta)$. The first case is trivial. In the second case we set $\zeta = a\mathbf{1} + z$. Since $\eta \in C'$ and $\eta \perp z$,

$$\begin{aligned} \zeta \eta \zeta &= (a\mathbf{1} + z)\eta(a\mathbf{1} + z) \\ &= a^2\eta + a(z\eta + \eta z) + z\eta z \\ &= a^2\eta + |z|^2\eta, \end{aligned}$$

which is in C .

3. For any v , $\Re(\zeta \cdot_v \eta) = \langle \zeta, \bar{\eta} \rangle$. In fact, both expressions are symmetric, bilinear, and they agree when $\zeta = \eta$.

Proposition 1.2. *Any division subalgebra of (C, \cdot_v) generated by two elements is associative.*

This is true in any normed division algebra. An easy classification-free proof is in [FK, pp. 82-83]. It is also well known and easy (cf. e.g. [FK]) that the only normed division algebras are $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$. But, for self-sufficiency of this article, here is a proof of the Proposition, making use of the present setup.

Proof. We look at the intersection of the subalgebra with C' . If it has dimension 0 or 1, the conclusion is trivial. If it has dimension at least 2, we can assume that the two generators z_1, z_2 are both in C' and orthonormal. Let $z_3 = z_1 \cdot_v z_2$. Then $z_3 \in C'$ and orthogonal to both z_1 and z_2 . In fact,

$$\begin{aligned} \langle z_3, \mathbf{1} \rangle |v|^2 &= \langle z_1 z_2 v, v \rangle = -\langle z_2 v, z_1 v \rangle = -\langle z_2, z_1 \rangle |v|^2 = 0, \\ \langle z_3, z_1 \rangle |v|^2 &= \langle z_1 z_2 v, z_1 v \rangle = \langle z_2 v, v \rangle = \langle z_2, \mathbf{1} \rangle |v|^2 = 0, \end{aligned}$$

and similarly $z_3 \perp z_2$. Furthermore,

$$(z_1 \cdot_v z_3)v = z_1(z_1 z_2 v) = z_1^2 z_2 v = -z_2 v,$$

so $z_1 \cdot_v z_3 = -z_2$. Similarly, $z_2 \cdot_v z_3 = z_1$. The Clifford relations imply that $z_i \cdot_v z_j = -z_j \cdot_v z_i$. This shows that $(C, \cdot_v) \cong \mathbb{H}$, which is associative. \square

Corollary 1.3. *If $\zeta, \eta, \lambda \in C$ belong to a division subalgebra of (C, \cdot_v) generated by two elements, and $\lambda \neq 0$, then*

$$\zeta \cdot_v \eta = \zeta \cdot_{\lambda v} \eta.$$

Proof. We have

$$\begin{aligned} (\zeta \cdot_{\lambda v} \eta)(\lambda v) &= \zeta \eta \lambda v = \zeta(\eta \cdot_v \lambda)v \\ &= (\zeta \cdot_v (\eta \cdot_v \lambda))v = ((\zeta \cdot_v \eta) \cdot_v \lambda)v \\ &= (\zeta \cdot_v \eta)(\lambda v), \end{aligned}$$

where we have used associativity of \cdot_v in the second line. Then use the fact that the map $\tau \mapsto \tau(\lambda v)$ from C to $C(\lambda v)$ is one-to-one. \square

Proposition 1.4. *If (C, V) is not irreducible, then the multiplication \cdot_v is independent of v .*

The proof we give is also contained in the proof of Theorem 1.1 of [CDKR1].

Proof. Given u, v non-zero elements of V , we prove that $\cdot_u = \cdot_v$. Assume first that $Cu + Cv$ is a direct sum. For $\zeta, \eta \in C$,

$$(\zeta \cdot_{u+v} \eta)(u + v) = (\zeta \cdot_{u+v} \eta)u + (\zeta \cdot_{u+v} \eta)v ,$$

but also

$$(\zeta \cdot_{u+v} \eta)(u + v) = \zeta \eta (u + v) = \zeta \eta u + \zeta \eta v = (\zeta \cdot_u \eta)u + (\zeta \cdot_v \eta)v .$$

Therefore $(\zeta \cdot_{u+v} \eta)u = (\zeta \cdot_u \eta)u$ and $(\zeta \cdot_{u+v} \eta)v = (\zeta \cdot_v \eta)v$, i.e. $\zeta \cdot_u \eta = \zeta \cdot_{u+v} \eta = \zeta \cdot_v \eta$.

Assume now that $Cu + Cv$ is not a direct sum. If $\zeta_0 u = \eta_0 v$ with $\zeta_0, \eta_0 \neq 0$, then $v = \eta_0^{-1} \zeta_0 u \in C(Cu) = Cu$, and so $Cv = Cu$. Since (C, V) is not irreducible, there is $v' \neq 0$ such that $v' \perp Cv$. Using the first part of the proof, $\cdot_v = \cdot_{v'} = \cdot_u$. \square

Corollary 1.5. *The product \cdot_v is independent of v if and only if (C, \cdot_v) is associative for one (and hence for all) v . This is the case when $\dim C \leq 4$.*

Proof. We have the general identity

$$(1.6) \quad \zeta \cdot_v (\eta \cdot_v \lambda) = (\zeta \cdot_{\lambda v} \eta) \cdot_v \lambda \quad (\forall \zeta, \eta, \lambda \in C) .$$

Then, if \cdot_v is independent of v , then (C, \cdot_v) is clearly associative. Assume now that, for some $v \neq 0$, (C, \cdot_v) is associative. By Proposition 1.4, we can restrict ourselves to the irreducible case $V = Cv$. So any other non-zero $u \in V$ is equal to λv for some non-zero $\lambda \in C$. By (1.6),

$$(\zeta \cdot_v \eta) \cdot_v \lambda = \zeta \cdot_v (\eta \cdot_v \lambda) = (\zeta \cdot_u \eta) \cdot_v \lambda .$$

Dividing by λ , we have $\zeta \cdot_v \eta = \zeta \cdot_u \eta$.

If $\dim C = 1$, then $C = \mathbb{R}\mathbf{1}$ and there is nothing to say. If $\dim C = 2$, taking $i \in C'$ with $i^2 = -1$, it is immediate to see that $C \cong \mathbb{C}$. If $\dim C \geq 3$, take orthonormal vectors $i, j \in C'$. Fix $v \in V$ a non-zero vector, and set $k = i \cdot_v j$. As shown in the proof of Proposition 1.2, k is linearly independent of $\mathbf{1}, i, j$, so that $\dim C = 4$ and (C, \cdot_v) is generated by i and j . By Proposition 1.2, $(C, \cdot_v) \cong \mathbb{H}$ is associative. \square

We shall say in short that C is associative, or non-associative, to distinguish between these two cases.

2. AUTOMORPHISMS OF $J^2 C$ -MODULES

Let (C, V) be a C -module. An automorphism of (C, V) is a pair $m = (\varphi, \psi)$ of orthogonal maps $\varphi : C \rightarrow C$, $\psi : V \rightarrow V$ such that the diagram

$$\begin{array}{ccc} C \times V & \xrightarrow{J} & V \\ \varphi \downarrow & \downarrow \psi & \downarrow \psi \\ C \times V & \xrightarrow{J} & V \end{array}$$

is commutative, i.e.

$$(2.1) \quad \psi(\zeta v) = \varphi(\zeta)\psi(v) .$$

We write M for the group of automorphisms of the C -module. The automorphism group of a general C -module is described in detail in [R]. We shall look more closely at the specific properties of M when V satisfies the J^2 -condition.

We denote by M_1 the subgroup of M defined by the condition $\varphi = \text{id}$. M_1 is the automorphism group of the Clifford module associated to (C, V) .

Proposition 2.1. *Let (C, V) be a $J^2 C$ -module, and assume that C is associative. Then M_1 is in one-to-one correspondence with the ordered orthonormal C -bases of V .*

Proof. Let $\{u_1, \dots, u_n\}$ and $\{v_1, \dots, v_n\}$ be two o.n. C -bases. Define $\psi : V \rightarrow V$ by

$$\psi\left(\sum \zeta_j u_j\right) = \sum \zeta_j v_j .$$

This is clearly a well-defined orthogonal map. To prove that $(\text{id}, \psi) \in M$, we must verify that

$$(2.2) \quad \psi(\eta v) = \eta \psi(v) \quad (\forall \eta \in C, v \in V) .$$

Since \cdot_v is independent of v ,

$$\begin{aligned} \psi\left(\eta \sum \zeta_j u_j\right) &= \psi\left(\sum (\eta \cdot_{u_j} \zeta_j) u_j\right) \\ &= \sum (\eta \cdot_{u_j} \zeta_j) v_j \\ &= \sum (\eta \cdot_{v_j} \zeta_j) v_j \\ &= \eta \sum \zeta_j v_j . \end{aligned}$$

It is then obvious that ψ is the only linear map satisfying (2.2) and mapping each u_j into v_j . \square

To construct automorphisms that are not trivial on C we need a preliminary remark about conjugation in $\text{Cliff}(C')$.

If z is a unit element of C' and $\eta \in C$, then the Clifford product $z\eta z^{-1} = -z\eta z$ is also in C (see Remark 2 in Section 1). In fact, it equals η if $\eta \in \mathbb{R}\mathbf{1} + \mathbb{R}z$, and $-\eta$ if $\eta \perp (\mathbb{R}\mathbf{1} + \mathbb{R}z)$ (in other words, $z\eta z^{-1}$ is the reflection of η in the plane $\mathbb{R}\mathbf{1} + \mathbb{R}z$).

Let $\text{Pin}(C') \subset \text{Cliff}(C')$ be the multiplicative group generated by unit elements of C' . It follows that, if $\alpha \in \text{Pin}(C')$, the map $\eta \mapsto \alpha\eta\alpha^{-1}$ is orthogonal on C and is the identity on $\mathbb{R}\mathbf{1}$.

On a general C -module, for any $\alpha \in \text{Pin}(C')$,

$$(2.3) \quad m_\alpha(\zeta, v) = (\alpha\zeta\alpha^{-1}, \alpha v) ,$$

is in M . The m_α form a subgroup that we denote by M_2 . M_2 acts on C' as $SO(C')$ if $\dim C'$ is odd, and as $O(C')$ if $\dim C'$ is even (this follows because this action is generated by the reflections $\eta \mapsto z\eta z^{-1}$). In particular, M_2 acts transitively on spheres in C' . It follows that M_2 is isomorphic with $\text{Pin}(C')$ modulo a finite subgroup. It is also true, even if of no great importance for us here, that $M_1 \cap M_2$ is finite and that the group $M_1 M_2$ has index at most 2 in M (cf. [R]).

Proposition 2.2. *Let (C, V) be a J^2C -module, and let Ξ_V be the manifold of ordered orthonormal C -bases of V . Let also $S_{C'}$ be the unit sphere in C' . Then M acts transitively on $S_{C'} \times \Xi_V$.*

Proof. For any $m = (\varphi, \psi) \in M$, $\varphi(\mathbf{1}) = \mathbf{1}$, hence φ acts orthogonally on C' . It is also simple to verify that ψ transforms orthonormal C -bases of V into orthonormal C -bases. Therefore M acts on $S_{C'} \times \Xi_V$.

Given two elements $(z_0, \xi_0), (z'_0, \xi'_0) \in S_{C'} \times \Xi_V$, we want to find $m \in M$ such that $m(z'_0, \xi'_0) = (z_0, \xi_0)$. There is $\alpha \in \text{Pin}(C')$ such that $\alpha z'_0 \alpha^{-1} = z_0$. Then α transforms ξ'_0 into another o.n. basis ξ'_1 . Then we need to find $m \in M$ such that $m(z_0, \xi'_1) = (z_0, \xi_0)$.

If C is associative, Proposition 2.1 says that there is an element of M_1 that does the job. If C is non-associative, we necessarily have $\dim C = \dim V \geq 5$, by Proposition 1.4 and Corollary 1.5. In particular, $\Xi_V = S_V$, the unit sphere in V , since (C, V) is irreducible. We then have two elements $v_1, v_2 \in S_V$, and we want to find $\beta \in \text{Pin}(C')$ such that $\beta z_0 \beta^{-1} = z_0$ and $\beta v_1 = v_2$.

Choose $v \in S_V$ orthogonal to v_j and $z_0 v_j$ for $j = 1, 2$. Since $V = Cv$, there are $z_1, z_2 \in C$ such that $v_j = z_j v$, $j = 1, 2$. By (1.4), $v \perp v_j$ implies that $z_j \in C'$, and $v \perp z_0 v_j$ implies that $z_j \perp z_0$. Let $\beta = -z_2 z_1 = z_2 z_1^{-1}$. Then

$$\begin{aligned} \beta z_0 \beta^{-1} &= z_2 z_1 z_0 z_1 z_2 = -z_2 z_0 z_2 = z_0, \\ \beta v_1 &= z_2 z_1^{-1} z_1 v = z_2 v = v_2. \quad \square \end{aligned}$$

Corollary 2.3. *M acts transitively on the product $S_{C'} \times S_V$ of the two unit spheres in C' and V , and on the product of unit spheres $S_{C'} \times S_{Cv_1} \times \cdots \times S_{Cv_n}$, if $\{v_1, \dots, v_n\}$ is an orthonormal C -basis of V .*

Transitivity of M on $S_{C'} \times S_V$ is *Kostant's double transitivity* for the $\text{Ad}(M)$ -action on the sum of root spaces $\mathfrak{g}^\lambda + \mathfrak{g}^{2\lambda}$ in semisimple Lie algebras.

3. CLASSIFICATION AND SOME BACKGROUND INFORMATION

Theorem 3.1 below gives the classification of all J^2C -modules in terms of normed division algebras. This theorem will not be used at all in the rest of the article.

Theorem 3.1. *Every J^2C -module is isomorphic with one of the following:*

- (i) C is any Euclidean space with a distinguished unit vector $\mathbf{1}$, $V = 0$;
- (ii) $C = \mathbb{F}$, a normed division algebra, $V = \mathbb{F}^n$ (with n a positive integer if \mathbb{F} is associative, and $n = 1$ if \mathbb{F} is non-associative, the norm on \mathbb{F}^n being the usual ℓ^2 -norm) and J is multiplication by elements of \mathbb{F} from the left.

Proof. These are clearly J^2C -modules. The J^2 -condition is vacuous in (i) and trivial in (ii) when \mathbb{F} is associative. If \mathbb{F} is non-associative, then $Cv = V$ for any $v \neq 0$, and the J^2 -condition follows trivially.

To prove that our list is complete, assume that (C, V) is a J^2C -module. If $V = 0$, there is nothing to prove. If $V \neq 0$, we must provide an isomorphism (f, g) from some $(\mathbb{F}, \mathbb{F}^n)$ as above to (C, V) , i.e. a pair of orthogonal linear maps $f : \mathbb{F} \rightarrow C$, $g : \mathbb{F}^n \rightarrow V$ such that

$$g(q(q_1, \dots, q_n)) = f(q)g(q_1, \dots, q_n) \quad (\forall q, q_1, \dots, q_n \in \mathbb{F}).$$

Fix an orthonormal C -basis $\{v_1, \dots, v_n\}$ of V and set $\mathbb{F} = (C, \cdot_{v_1})$. Let $f : \mathbb{F} \rightarrow C$ be the identity map and define $g : \mathbb{F}^n \rightarrow V$ by

$$g(\zeta_1, \dots, \zeta_n) = \sum \zeta_j v_j .$$

If \mathbb{F} is associative, $\cdot_{v_1} = \cdot_{v_j}$ for every j . Then (as in the proof of Proposition 2.1)

$$\begin{aligned} g(\eta \cdot_{v_1} (\zeta_1, \dots, \zeta_n)) &= \sum (\eta \cdot_{v_1} \zeta_j) v_j \\ &= \sum \eta \zeta_j v_j \\ &= \eta \sum \zeta_j v_j \\ &= f(\eta) g(\zeta_1, \dots, \zeta_n) . \end{aligned}$$

This computation also works when C is non-associative, due to the fact that $n = 1$. \square

We conclude this section by mentioning the relations among C -modules, Clifford modules and H -type Lie algebras.

As pointed out in Section 1, every C -module extends in a natural way to a $\text{Cliff}(C')$ -module. This construction can be reversed: let C' be any finite dimensional (possibly trivial) vector space with a scalar product and V a module over $\text{Cliff}(C')$, endowed with a $\text{Pin}(C')$ -invariant scalar product. Setting $C = \mathbb{R}\mathbf{1} \oplus C'$, there is a unique C -module structure (C, V) inducing the given Clifford module structure on V [K]. However, non-equivalent Clifford modules may induce isomorphic C -modules. This is because (C, V) and (C, V') may be isomorphic without any isomorphism being the identity on C (cf. [KR]).

The main facts about Clifford modules can be derived from [Hus]. If the dimension d of C is not divisible by 4, $\text{Cliff}(C')$ has, up to equivalence, only one irreducible module V_0 , and every other module can be realized as $V_k \sim V_0 \otimes_{\mathbb{F}} \mathbb{F}^k$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$, depending on the congruence class of $d \pmod{8}$. If d is divisible by 4, then $\text{Cliff}(C')$ has two inequivalent irreducible modules, V_1, V_2 , and any module V can be realized as $V_{kh} = (V_1 \otimes_{\mathbb{F}} \mathbb{F}^k) \oplus (V_2 \otimes_{\mathbb{F}} \mathbb{F}^h)$, with $\mathbb{F} = \mathbb{R}$ if d is divisible by 8, and $\mathbb{F} = \mathbb{H}$ otherwise. If $k \neq h$, V_{kh} and V_{hk} are non-equivalent as Clifford modules, but the induced C -modules are isomorphic.

The J^2 -condition makes sense on Clifford modules, and it can be restated as $\text{Cliff}(C')v = Cv$ for every $v \in V$. This can only occur when $\text{Cliff}(C')$ has a module (necessarily irreducible) of the same dimension as C , i.e. when $d = 1, 2, 4, 8$. For these values of d , the J^2 -condition holds for every module if $d = 1, 2$, only for the “isotypic” C -modules $V_{k0} = V_{0k}$ if $d = 4$, and only for the irreducible C -module if $d = 8$ [CDKR1].

There is a third category which must be mentioned in this context: the H -type Lie algebras. An H -type Lie algebra (the notion of which is the basic object in [CDKR1,2]) is a Lie algebra \mathfrak{n} with a positive definite inner product. It is assumed that \mathfrak{z} is the center of \mathfrak{n} , $\mathfrak{n} = \mathfrak{v} + \mathfrak{z}$ is an orthogonal direct sum and $[\mathfrak{v}, \mathfrak{v}] \subset \mathfrak{z}$. Furthermore, for all $z \in \mathfrak{z}$, the map $J_z : \mathfrak{v} \rightarrow \mathfrak{v}$, defined by

$$(3.1) \quad \langle J_z v, u \rangle = \langle [v, u], z \rangle ,$$

has the property

$$(3.2) \quad J_z^2 = -|z|^2 \text{id}_{\mathfrak{v}} .$$

It is immediate that, given a C -module (C, V) , we obtain an H -type Lie algebra by taking \mathfrak{n} as the Euclidean direct sum of $\mathfrak{v} = V$ and $\mathfrak{z} = C'$ and defining the Lie bracket through (3.1). The converse is also easy to verify, as well as that isomorphic C -modules produce isomorphic Lie algebras, and *vice versa* [K, KR].

4. THE SPACE $W = C \oplus V$

Given a J^2C -module (C, V) , let $W = C \oplus V$ be the direct sum of Euclidean spaces. We introduce the following equivalence relation on $W \setminus \{0\}$:

- (i) $(0, u) \sim (0, v)$ if $u \in Cv$;
- (ii) if $\zeta \neq 0$, $(\eta, u) \sim (\zeta, v)$ if $\eta \neq 0$ and $\eta^{-1}u = \zeta^{-1}v$.

Notice that the J^2 -condition is required in (i) to prove transitivity. Given an element $w = (\zeta, v) \neq 0$ in W , we denote its equivalence class together with point 0 by Cw or $C(\zeta, v)$. We must pay attention, however, to the fact that the notation ηw does not make any sense for an individual $\eta \in C$. This is related to the fact that in general W cannot be given a C -module structure in a natural way (unless C is associative). We only have that, for $v \neq 0$, $C(\zeta, v) = \{(\lambda \cdot_v \zeta, \lambda v) : \lambda \in C\} = \{(\lambda, \lambda \zeta^{-1}v) : \lambda \in C\}$.

We call C -line, or *affine C -line*, any translate $w_0 + Cw$ of Cw , with $w \neq 0$. The C -lines through 0 form a closed subset of the (Grassmanian) manifold of real d -dimensional subspaces of W . To see this, one has to show that if $(\zeta_n, v_n) \rightarrow (\zeta, v)$, $(\zeta'_n, v'_n) \rightarrow (\zeta', v')$ and $(\zeta_n, v_n) \sim (\zeta'_n, v'_n)$, then $(\zeta, v) \sim (\zeta', v')$. This is easy and left to the reader. Since different C -lines through 0 meet only at 0, we also see that $w_n \rightarrow w \neq 0$ implies that $Cw_n \rightarrow Cw$.

It is clear that every C -line through 0 can be written in the form $C(\mathbf{1}, v)$ (uniquely), or $C(0, v)$ (non-uniquely). The C -lines of the first type form a dense open set (dense because $C(0, v) = \lim C(\frac{1}{n}, v) = \lim_{n \rightarrow \infty} C(\mathbf{1}, nv)$ as $n \rightarrow \infty$).

We call $GL(W, C)$ the group of \mathbb{R} -linear transformations of W preserving C -lines. We also set $K = GL(W, C) \cap O(W)$. Since $GL(W, C)$ and K are closed subgroups of $GL(W, \mathbb{R})$, they are Lie groups.

Proposition 4.1. *If $(\varphi, \psi) \in M$, the map $\varphi \times \psi : W \rightarrow W$ defined by $(\varphi \times \psi)(\zeta, v) = (\varphi(\zeta), \psi(v))$ is in K . This correspondence identifies M with the subgroup of K whose elements fix the point $(\mathbf{1}, 0)$. The subgroup M_1 of M is then identified with the subgroup of K whose elements fix every point in C .*

Proof. If $(\varphi, \psi) \in M$, then $\varphi(\mathbf{1}) = \mathbf{1}$, so that $\varphi \times \psi$ fixes $(\mathbf{1}, 0)$. Let $\ell = C(\zeta, v)$ be a C -line through 0. If $\zeta = 0$, then $\ell = \{(0, \eta v) : \eta \in C\}$, and

$$(\varphi \times \psi)(0, \eta v) = (0, \psi(\eta v)) = (0, \varphi(\eta)\psi(v)) \in C(\varphi \times \psi)(0, v) .$$

If $\zeta \neq 0$, then $\ell = C(\mathbf{1}, \zeta^{-1}v)$, so we may as well assume that $\zeta = \mathbf{1}$. Then $\ell = \{(\eta, \eta v) : \eta \in C\}$ and

$$(\varphi \times \psi)(\eta, \eta v) = (\varphi(\eta), \varphi(\eta)\psi(v)) \in C(\mathbf{1}, \psi(v)) = C(\varphi \times \psi)(\mathbf{1}, v) .$$

Conversely, assume that $k \in K$ fixes $(\mathbf{1}, 0)$. Then kC is the C -line containing $(\mathbf{1}, 0)$, i.e. C . By orthogonality, $kV = V$. Denote by k_1, k_2 the restrictions of k to C and V respectively. Consider now an element $(\eta, \eta v)$ in a C -line $C(\mathbf{1}, v)$ not contained in V . By linearity, $k(\eta, \eta v) = (k_1(\eta), k_2(\eta v))$. Moreover, $k(\eta, \eta v) \in Ck(\mathbf{1}, v) = C(\mathbf{1}, k_2(v))$. Then necessarily $k_2(\eta v) = k_1(\eta)k_2(v)$, i.e. $(k_1, k_2) \in M$.

The last part of the statement is now obvious. \square

We will construct elements of K that are not in M , i.e. that do not fix $(\mathbf{1}, 0)$. But first we prove a more general statement, which will also be useful later.

Proposition 4.2. *Let $\{v_1, \dots, v_n\}$ be an orthonormal C -basis of V . Let $w_0 = (\mathbf{1}, 0)$, $w_j = (0, v_j)$ ($1 \leq j \leq n$) and $(a_{jk})_{0 \leq j, k \leq n}$ be a real matrix. Then the map $A : W \rightarrow W$ defined by*

$$A\left(\sum_{j=0}^n \zeta_j w_j\right) = \sum_{j,k=0}^n a_{jk} \zeta_k w_j$$

is in $GL(W, C)$.

Though the expression ζw does not make sense in general, the abuse of notation in the statement above does not cause ambiguity. It must be understood that $\zeta_0 w_0 = (\zeta_0, 0)$ and $\zeta_j w_j = (0, \zeta_j v_j)$ for $j \geq 1$.

Proof. It is clear that A is well defined and \mathbb{R} -linear, so that it suffices to prove the last statement for C -lines through 0. By continuity, we can also restrict ourselves to C -lines $\ell = C(\mathbf{1}, v)$. Now, for $v = \sum_{j=1}^n \zeta_j v_j$,

$$C(\mathbf{1}, v) = \left\{ \left(\lambda, \sum_{j=1}^n \lambda \zeta_j v_j \right) : \lambda \in C \right\} = \left\{ \sum_{j=0}^n (\lambda \cdot_{v_j} \zeta_j) w_j : \lambda \in C \right\}$$

(where $\zeta_0 = \mathbf{1}$ and $\lambda \cdot_{v_0} \zeta_0$ stands for λ). So,

$$A(\ell) = \left\{ \sum_{j,k=0}^n a_{jk} (\lambda \cdot_{v_k} \zeta_k) w_j : \lambda \in C \right\} .$$

This is clearly a C -line if C is associative, because \cdot_{v_k} is independent of k .

If C is non-associative, then $n = 1$ and we must verify that

$$\begin{aligned} & (a_{00}\lambda + a_{01}\lambda \cdot_{v_1} \zeta_1)^{-1} (a_{10}\lambda + a_{11}\lambda \cdot_{v_1} \zeta_1) \\ &= \left[(a_{00}\mathbf{1} + a_{01}\zeta_1)^{-1} \cdot_{v_1} \lambda^{-1} \right] \cdot_{(a_{10}\lambda + a_{11}\lambda \cdot_{v_1} \zeta_1)v_1} \left[\lambda \cdot_{v_1} (a_{10}\mathbf{1} + a_{11}\zeta_1) \right] \end{aligned}$$

does not depend on λ . For fixed λ , every element of C appearing in this expression belongs to the division subalgebra of (C, \cdot_{v_1}) generated by ζ_1 and λ . By Proposition 1.2 and Corollary 1.3, this subalgebra is associative and $\cdot_{(a_{10}\lambda + a_{11}\lambda \cdot_{v_1} \zeta_1)v_1} = \cdot_{v_1}$, so that λ cancels out. \square

Proposition 4.3. *For a fixed unit vector $v_0 \in V$, decompose $v \in V$ as $v = \eta v_0 + v'$, with $v' \in (Cv_0)^\perp$, and set, for $\theta \in \mathbb{T}$,*

$$\sigma_{v_0, \theta}(\zeta, \eta v_0 + v') = (\cos \theta \zeta - \sin \theta \eta, (\cos \theta \eta + \sin \theta \zeta)v_0 + v') .$$

Then $\sigma_{v_0, \theta} \in K$.

Proof. It is trivial to verify that $\sigma_{v_0, \theta} \in O(W)$. It maps C -lines into C -lines because v_0 can be completed to an o.n. basis of V and then Proposition 4.2 applies. \square

Corollary 4.4. *K acts transitively on the unit sphere S_W (and hence on C -lines through 0).*

Proof. Given a point $(\zeta, v) \in S_W$, write (ζ, v) as $(c\zeta', sv')$ with $\zeta' \in S_C$, $v' \in S_V$, $c = \cos \theta$, $s = \sin \theta$ for an appropriate $\theta \in \mathbb{T}$ (if $v = 0$, we choose v' arbitrarily). Then $k = \sigma_{\zeta'^{-1}v', \theta - \frac{\pi}{2}} \circ \sigma_{v', \frac{\pi}{2}}$ maps $(\mathbf{1}, 0)$ into (ζ, v) . \square

We call *linear C -subspace* of W an \mathbb{R} -linear subspace E such that $Cw \subset E$ whenever $w \neq 0$ is in E .

We also call *affine C -subspace* of W a translate $E' = E + w$ of a linear C -subspace E . This is equivalent to saying that E' is a set that contains the whole affine C -line $w_1 + C(w_2 - w_1)$ connecting any pair of points $w_1, w_2 \in E'$.

For an \mathbb{R} -linear subspace of V , to be a sub- C -module of V is the same as being a linear C -subspace of W .

Lemma 4.5. *The orthogonal of a linear C -subspace is a linear C -subspace.*

Proof. Let E be a linear C -subspace. Modulo the action of K , we can assume that $C \subset E$. Then $E = C \oplus V_0$, with V_0 a sub-module of V . Then the orthogonal of E in W is the same as the orthogonal of V_0 in V , which is also a sub-module. \square

Proposition 4.6. *The linear span in W of C -lines through 0 is a linear C -subspace. Conversely, any linear C -subspace is the orthogonal sum of C -lines through 0.*

Proof. The first part of the statement follows by induction, if we prove that the sum of a proper linear C -subspace E and a C -line ℓ through 0 is a linear C -subspace.

Modulo the action of K , we can assume that E^\perp contains C , hence that $E \subset V$. If ℓ is also contained in V , then $E + \ell$ is a sub-module of V .

If $\ell \not\subset V$, then $\ell = C(\mathbf{1}, v_0)$ for some $v_0 \in V$ and

$$E + \ell = \{(\eta, \eta v_0 + v) : \eta \in C, v \in S\} .$$

We need to prove that, if $\eta \neq 0$ and $v \in E$, then $C(\eta, \eta v_0 + v) \subset E + \ell$. An element of $C(\eta, \eta v_0 + v)$ has the form $(\zeta, \zeta v_0 + \zeta \eta^{-1}v)$, which is in $E + \ell$, being $\zeta \eta^{-1}v \in E$.

For the converse, we can assume that $C \subset E$. Then $E = C \oplus V_0$, with V_0 a sub- C -module of V , and it can be decomposed into the orthogonal sum of irreducible ones. \square

Proposition 4.6 allows us to introduce the notions of *orthonormal C -basis* and of *C -dimension* of a linear C -subspace of W . We can then extend the scope of Corollary 4.4.

Corollary 4.7. *The group K acts transitively on the manifold Ξ_W of ordered orthonormal C -bases of W and on the manifold of linear C -subspaces of any fixed dimension.*

Proof. It follows from Proposition 2.2 that $M \subset K$ acts transitively on ordered orthonormal C -bases of W whose first element is $(\mathbf{1}, 0)$. It is then sufficient to prove that any other basis $\{w_0, \dots, w_n\}$ can be mapped into a basis of this type. But this amounts to saying that there is $k \in K$ such that $kw_0 = (\mathbf{1}, 0)$, and this follows from Corollary 4.4. \square

In the remainder of this section we describe the orbits of M on the unit sphere S_W of W and analyze the structure of K in more detail.

We assume that $C' \neq 0$ and $V \neq 0$, the degenerate cases being trivial. We fix an o.n. C -basis $\{v_1, \dots, v_n\}$ of V and an element $z \in C'$ with $|z| = 1$.

Lemma 4.8. *Every M -orbit in S_W meets the subspace $\mathbb{R}\mathbf{1} + \mathbb{R}z + \mathbb{R}v_1$ in the points $a\mathbf{1} \pm bz \pm cv_1$ for unique numbers $a \in \mathbb{R}$, $b, c \geq 0$ with $a^2 + b^2 + c^2 = 1$.*

Proof. M fixes $\mathbf{1}$, hence $\Re \zeta$ remains constant on any orbit $M \cdot (\zeta, v)$. The rest of the statement follows from Corollary 2.3. \square

We write, for $t \in \mathbb{R}$, $\eta_t = (\cos t)\mathbf{1} + (\sin t)z$, and define m_t by

$$m_t(\zeta, v) = (\eta_t \zeta \eta_t^{-1}, \eta_t v) .$$

This is a one-parameter subgroup of M ($m_t \in M_2$ if $\dim C' > 1$, and $m_t \in M_1$ if $\dim C' = 1$). We write $\sigma_1 = \sigma_{v_1, \frac{\pi}{2}}$ and define T as the one-parameter subgroup of K consisting of the elements

$$\rho_t = \sigma_1^{-1} \circ m_t \circ \sigma_1 .$$

We also denote by L the subgroup of K preserving C (hence also preserving V). Then $M \subset L$, and also $T \subset L$, since σ_1 only interchanges C with Cv_1 and preserves every other Cv_j .

For any $v \in V$ and $t \in \mathbb{R}$, we have $\rho_t(\mathbf{1}, v) = (z_t, v')$ for some $v' \in V$. Hence the orbits of L on S_W all meet the plane $\mathbb{R}\mathbf{1} + \mathbb{R}v_1$ (in points $\pm a\mathbf{1} \pm cv_1$, for unique $a, c \geq 0$ with $a^2 + c^2 = 1$).

We write T' for the subgroup $\{\sigma_{v_1, \theta}\}_{\theta \in \mathbb{T}}$ of K .

Proposition 4.9. *We have $K = LT'L$ and $L = MTM$.*

The proof is trivial from the structure of the L - and M -orbits in S_W .

Corollary 4.10. *The following hold:*

- (i) M and T generate K ;
- (ii) $MkM = Mk^{-1}M$ for every $k \in K$;
- (iii) the projection of the action of L onto C is $SO(C)$ or $O(C)^1$.

Proof. (i) is obvious and (ii) follows from Lemma 4.8. As to (iii), the description of M given in Section 2 shows that M_2 acts on C as $SO(C')$ or $O(C')$. On the other hand, T acts on C as a torus transversal to $SO(C')$. It is well known, and easy, that $SO(d-1)$ is a maximal proper subgroup of $SO(d)$, and the conclusion follows. \square

Remark 1. For future reference, we observe that, given $w, w' \in W$ and writing π_{Cw} for the orthogonal projection on Cw , the ratio $|\pi_{Cw} w''|/|w''|$ is constant for all $w'' \in Cw'$. This constant is then the cosine of the angle of Cw and Cw' . The proof is immediate when $Cw = C$. The general case follows by Corollary 4.7.

Remark 2. Corollary 4.10(ii) implies that (K, M) is a Gelfand pair. This and (iii) are the key facts in the discussion of harmonic analysis on K/M in the exceptional case developed in [Ta]. Here we have proved them without explicit use of the octonions.

¹The projection of the action of L onto C is $SO(C)$ if $d = 4, 8$ and $O(C)$ if $d = 1, 2$.

5. THE CONSTRUCTION OF CPW

The compact rank-one symmetric spaces will be defined as appropriate compactifications of the vector spaces W that we have associated to J^2C -modules. In the associative case, these compactifications can be described (as they are in the literature) as spaces of lines: according to Theorem 3.1, we identify C with an associative division algebra \mathbb{F} , V with \mathbb{F}^n and W with \mathbb{F}^{n+1} . The symmetric space is then $\mathbb{F}P^{n+1} = (\mathbb{F}^{n+2} \setminus \{0\})/\mathbb{F}^*$. This construction makes use of the fact that we can “add dimensions” to V without destroying the J^2C -module structure, something that cannot be adapted to the non-associative case.

In order to have a unified description, including both the associative and the non-associative case, we construct the compactification by “gluing” to W a space W_∞ consisting of “points at infinity”.

We set $W_\infty = (W \setminus \{0\})/\sim$, where \sim is the equivalence relation introduced at the beginning of Section 4. It is convenient to think of W_∞ as the set of C -lines through 0 in W . We denote by $\pi : W \setminus \{0\} \rightarrow W_\infty$ the quotient map, and we endow W_∞ with the quotient topology. The element $\pi(w)$ will be denoted by $[w]$.

We then set $CPW = W \cup W_\infty$, and define a topology on it by assigning neighborhood bases at the various points as follows:

- (i) to a point $w \in W$, we assign its Euclidean neighborhoods in W ;
- (ii) to a point $[w] \in W_\infty$, we assign, for each neighborhood $U_{[w]}^\infty$ of $[w]$ in W_∞ and each $R > 0$, the neighborhood

$$U_{[w]}^R = U_{[w]}^\infty \cup \{w' \in \pi^{-1}(U_{[w]}^\infty) : |w'| > R\} .$$

We leave the reader the verification that these neighborhood systems actually define a second-countable topology. It is obvious that W is open and dense in CPW .

As we have seen, a C -line not contained in V contains a unique element $(\mathbf{1}, v)$. We denote the corresponding element of W_∞ as $[\mathbf{1}, v]$ and W_∞^1 the set of such elements. The complementary subset W_∞^0 of W_∞ is the set of C -lines contained in V . We use the notation $[0, v]$ to denote the C -line containing $v \neq 0$ (in this case, however, $[0, v]$ depends only on the equivalence class of v mod. C^*). Clearly, W_∞^1 is open and dense in W_∞ .

The following facts about convergence of sequences in CPW are easily derived from the definition:

- (i) $W \ni w_n \rightarrow [w] \in W_\infty$ if and only if $|w_n| \rightarrow \infty$ and $[w_n] \rightarrow [w]$ in W_∞ ;
- (ii) $(\zeta_n, v_n) \rightarrow [\mathbf{1}, v] \in W_\infty^1$ if and only if $|\zeta_n| \rightarrow \infty$ and $\zeta_n^{-1}v_n \rightarrow v$;
- (iii) $(\zeta_n, v_n) \rightarrow [0, v] \in W_\infty^0$ if and only if $|v_n| \rightarrow \infty$, $|\zeta_n| = o(|v_n|)$ and $\exists \lambda_n (\rightarrow 0)$ in C such that $\lambda_n v_n \rightarrow v$;
- (iv) $W_\infty^1 \ni [\mathbf{1}, v_n] \rightarrow [0, v] \in W_\infty^0$ if and only if $|v_n| \rightarrow \infty$ and $\exists \lambda_n (\rightarrow 0)$ such that $\lambda_n v_n \rightarrow v$.

We state a couple of facts that will be used later on.

Proposition 5.1. *Let $f \in GL(W, C)$. The induced map $f_\infty : W_\infty \rightarrow W_\infty$ such that $f_\infty \circ \pi = \pi \circ f$ is a homeomorphism and $\bar{f} = f \cup f_\infty$ is a homeomorphism of CPW .*

Translations in W extend to homeomorphisms of CPW which are the identity on W_∞ .

Proof. That f_∞ is a homeomorphism follows from standard facts about quotient topologies. It is then sufficient to prove that $f(w_n) \rightarrow f_\infty([w])$ if $W \ni w_n \rightarrow [w] \in W_\infty$. Using (i) above, $|f(w_n)| \rightarrow \infty$ because f is invertible, and

$$[f(w_n)] = f_\infty([w_n]) \rightarrow f_\infty([w]) .$$

Take now a translation $\tau(w) = w + w_0$ with $w_0 = (\zeta_0, v_0)$, and assume that $(\eta_n, u_n) \rightarrow [w] \in W_\infty$. We claim that also $(\eta_n + \zeta_0, u_n + v_0) \rightarrow [w]$.

If $[w] = [\mathbf{1}, v] \in W_\infty^1$, by (ii) we are assuming that $|\eta_n| \rightarrow \infty$ and $v_n = \eta_n^{-1}u_n \rightarrow v$. Then $|\eta_n + \zeta_0| \rightarrow \infty$ and

$$\begin{aligned} \lim_{n \rightarrow \infty} (\eta_n + \zeta_0)^{-1}(u_n + v_0) &= \lim_{n \rightarrow \infty} (\eta_n + \zeta_0)^{-1}u_n \\ &= \lim_{n \rightarrow \infty} (\eta_n + \zeta_0)^{-1}\eta_n v_n \\ &= v - \lim_{n \rightarrow \infty} (\eta_n + \zeta_0)^{-1}\zeta_0 v_n \\ &= v . \end{aligned}$$

If $[w] = [0, v] \in W_\infty^0$, by (iii) we are assuming that $|u_n| \rightarrow \infty$, $|\eta_n| = o(|u_n|)$ and $\lambda_n u_n \rightarrow v$ for some sequence $\lambda_n \rightarrow 0$. Then $|u_n + v_0| \rightarrow \infty$, $|\eta_n + \zeta_0| \sim |\eta_n| = o(|u_n + v_0|)$, and $\lambda_n(u_n + v_0) \rightarrow v$. \square

From now on we will use the same symbol for a C -linear or C -affine map of W and for its continuous extension to CPW .

Proposition 5.2. *If E is a linear C -subspace, the closure in CPW of the affine C -subspace $E' = w_0 + E$ is $(w_0 + E) \cup \pi(E)$.*

Proof. By Proposition 5.1, we can assume that $w_0 = 0$. Since $\pi(E) = \pi(E \cap S_W)$, $\pi(E)$ is closed in W_∞ . By (i), if $w_n \rightarrow [w]$ and $w_n \in E$, then $[w_n] \rightarrow [w]$, hence $[w] \in \pi(E)$. Conversely, if $[w] \in \pi(E)$, then $nw \rightarrow [w]$. \square

In order to introduce a differentiable structure on CPW compatible with this topology, we first show that certain maps on CPW are homeomorphisms. These maps will then be used to define the coordinate patches.

Lemma 5.3. *Let $\varphi_0 : CPW \rightarrow CPW$ be defined as*

$$(5.1) \quad \begin{aligned} \varphi_0(\zeta, v) &= (\zeta^{-1}, \zeta^{-1}v) \quad \text{if } \zeta \neq 0 \\ \varphi_0(0, v) &= [\mathbf{1}, v] \\ \varphi_0[\mathbf{1}, v] &= (0, v) \\ \varphi_0[0, v] &= [0, v] . \end{aligned}$$

Then φ_0 is a homeomorphism.

Proof. Since $\varphi_0^{-1} = \varphi_0$, it is sufficient to prove that φ_0 is continuous. The following facts must be verified:

- (1) $(\zeta_n, v_n) \rightarrow (0, v)$ (with $\zeta_n \neq 0$) if and only if $(\zeta_n^{-1}, \zeta_n^{-1}v_n) \rightarrow [\mathbf{1}, v]$;
- (2) if $(\zeta_n, v_n) \rightarrow [0, v]$ (with $\zeta_n \neq 0$), then $(\zeta_n^{-1}, \zeta_n^{-1}v_n) \rightarrow [0, v]$;
- (3) $(0, v_n) \rightarrow [0, v]$ if and only if $[\mathbf{1}, v_n] \rightarrow [0, v]$.

These can be easily verified on the basis of (ii)-(iv). \square

Fix now an orthonormal C -basis $\{v_1, \dots, v_n\}$ of V , and define, for $j = 1, \dots, n$ and $v' \perp Cv_j$,

$$\psi_j(\zeta, \eta v_j + v') = (\eta, \zeta v_j + v') .$$

Then ψ_j is linear and involutive on W and maps C -lines into C -lines. Therefore it extends to an involutive homeomorphism, also denoted by ψ_j , of CPW . If $\varphi_j = \varphi_0 \circ \psi_j \circ \varphi_0$, then also φ_j is an involutive homeomorphism of CPW . Explicitly, with $v' \perp Cv_j$,

$$(5.2) \quad \begin{aligned} \varphi_j(\zeta, \eta v_j + v') &= (\eta^{-1} \cdot_{v_j} \zeta, \eta^{-1} v_j + \eta^{-1} v') && \text{if } \eta \neq 0 \\ \varphi_j(\zeta, v') &= \begin{cases} [\mathbf{1}, \zeta^{-1}(v_j + v')] & \text{if } \zeta \neq 0, \\ [0, v_j + v'] & \text{if } \zeta = 0, \end{cases} \\ \varphi_j[\mathbf{1}, \eta v_j + v'] &= \begin{cases} (\eta^{-1}, \eta^{-1} v') & \text{if } \eta \neq 0, \\ [\mathbf{1}, v'] & \text{if } \eta = 0, \end{cases} \\ \varphi_j[0, \eta v_j + v'] &= \begin{cases} (0, \eta^{-1} v') & \text{if } \eta \neq 0, \\ [0, v'] & \text{if } \eta = 0. \end{cases} \end{aligned}$$

For notational convenience, we set $\varphi_{n+1} = \text{id}_{CPW}$.

Proposition 5.4. *The charts $(\varphi_j(W), \varphi_j)$, with $0 \leq j \leq n+1$, define a differentiable structure on CPW .*

(Since $\varphi_j = \varphi_j^{-1}$, the coordinate maps take values in W .)

Proof. We show that $W, \varphi_0(W), \dots, \varphi_n(W)$ form an open covering of CPW . In fact, the points of W_∞ that are not contained in $\varphi_0(W)$ are those in $\pi(V)$, whereas, for $j \geq 1$, the points of W_∞ not contained in $\varphi_j(W)$ are those in $\pi((Cv_j)^\perp)$. Since

$$V \cap (Cv_1)^\perp \cap \dots \cap (Cv_n)^\perp = 0 ,$$

we conclude that

$$\bigcup_{j=0}^n \varphi_j(W) \supset W_\infty .$$

We see directly from (5.1) and (5.2) that each component of φ_j , $0 \leq j \leq n$, is a rational function. The same is then true also for the transition maps $\varphi_j \circ \varphi_k$. In particular they are smooth (and analytic) on $W \cap (\varphi_j \circ \varphi_k)^{-1}W$. \square

When restricted to W_∞ , this differentiable structure coincides with the quotient structure of $(W \setminus \{0\})/\sim$. Restricting the quotient map π to the unit sphere S_W , we obtain the *Hopf fibration*

$$\pi : S_W \rightarrow W_\infty ,$$

with fiber S_C . By Proposition 5.1, K acts by diffeomorphisms of W_∞ and, by Corollary 4.7, this action is transitive. Let L be the subgroup of K introduced in Section 4, whose elements preserve C as a set. Identifying W_∞ with K/L and the unit sphere S_W with K/M , it follows that L is the structure group of the bundle, and the fiber S_C is diffeomorphic to L/M .

It may help the reader to see explicitly how CPW identifies with $\mathbb{F}P^{n+1}$, when \mathbb{F} is an associative division algebra, $C = \mathbb{F}$ and $V = \mathbb{F}^n$. Let $\Pi : (\mathbb{F}^{n+2} \setminus \{0\}) \rightarrow \mathbb{F}P^{n+1}$

be the quotient map. We think of W inside CPW as $\Pi(\mathbb{F}^{n+1} \times \{1\})$ and of W_∞ as $\Pi(\mathbb{F}^{n+1} \times \{0\})$. Then each φ_j is the projective image of the map interchanging the j -th component of $(q_0, q_1, \dots, q_n, q_{n+1}) \in \mathbb{F}^{n+2}$ with the $(n+1)$ -th component.

6. CPW AS A COMPACT SYMMETRIC SPACE

We first introduce a metric on W and then prove that it can be extended to all of CPW . Guided by the non-compact case (cf. (A.1) in the Appendix), we set

$$(6.1) \quad \langle X, Y \rangle_{w+} = \begin{cases} \frac{\langle X, Y \rangle}{(1 + |w|^2)^2} & \text{if } X, Y \in Cw, \\ \frac{\langle X, Y \rangle}{1 + |w|^2} & \text{if } X, Y \in (Cw)^\perp, \\ 0 & \text{if } X \in Cw, Y \in (Cw)^\perp, \end{cases}$$

if $w \neq 0$ and, for $w = 0$,

$$(6.2) \quad \langle X, Y \rangle_{0+} = \langle X, Y \rangle \quad \text{for every } X, Y.$$

Notice that (6.1) and (6.2) imply that for arbitrary $X \in W$,

$$(6.3) \quad |X|_{w+} = c_{|w|, \varphi} |X|,$$

where φ is the angle of the C -lines Cw and CX (see Remark 1 at the end of Section 4) and

$$(6.4) \quad c_{|w|, \varphi} = \left(\frac{\cos^2 \varphi}{(1 + |w|^2)^2} + \frac{\sin^2 \varphi}{1 + |w|^2} \right)^{\frac{1}{2}}.$$

Proposition 6.1. *The geodesic γ going through the origin with tangent vector $w \in S_W$ is $\gamma(t) = (\tan t)w$, $|t| < \frac{\pi}{2}$. The group of isometries of W fixing the origin is K .*

Proof. The elements of K are isometries of W , as a direct consequence of their property of mapping C -lines into C -lines. We can then assume that $w = (\mathbf{1}, 0)$. Then γ must be invariant under M . By Kostant's double transitivity (Corollary 2.3), this implies that $\gamma \subset \mathbb{R}\mathbf{1}$, i.e. $\gamma(t) = r(t)w$. We impose now that

$$\|\dot{\gamma}(t)\|_{\gamma(t)+} = \frac{r'(t)}{1 + r^2(t)} = 1.$$

This gives $r(t) = \tan t$.

Finally, let f be an isometry of W with $f(0) = 0$, and let $k = Df(0) \in O(W)$. If $w \in S_W$, the geodesic $\gamma(t) = (\tan t)w$ is mapped into $f(\gamma(t)) = (\tan t)k(w) = k(\gamma(t))$. Therefore $f = k$. Then (6.1) implies that k must map C -lines into C -lines. \square

We will see that the transition maps $\varphi_j \circ \varphi_k$ are isometric on $W \cap (\varphi_j \circ \varphi_k)^{-1}(W)$, and this will allow us to extend the metric to CPW , defining it locally as the pull-back of the metric in W via the coordinate maps φ_j . Since $\psi_j \in K$ for $1 \leq j \leq n$, the crucial fact to be proved is that φ_0 is an isometry on $W \setminus V$.

This will be the first consequence of the next lemma.

Lemma 6.2. For $\theta \in \mathbb{T}$ and $\zeta \neq (\cot \theta)\mathbf{1}$, define

$$(6.5) \quad b_\theta(\zeta, v) = ((\cos \theta \mathbf{1} - \sin \theta \zeta)^{-1}(\sin \theta \mathbf{1} + \cos \theta \zeta), (\cos \theta \mathbf{1} - \sin \theta \zeta)^{-1}v) ,$$

where the product in the first component is in the sense of Clifford multiplication or, equivalently, with respect to any product \cdot_{v_0} . Then b_θ is an isometry.

Before giving the proof, we comment that the b_θ must be regarded as close analogues of the maps $\sigma_{v_0, \theta}$ introduced in Proposition 4.3. We can see this in the associative case, where $CPW = \mathbb{F}P^{n+1}$. In terms of the homogeneous coordinates (q_0, \dots, q_{n+1}) introduced at the end of the previous section, and with v_0 the j -th basis element ($1 \leq j \leq n$), $\sigma_{v_0, \theta}$ introduces a rotation by θ in the pair of coordinates (q_0, q_j) . Now b_θ introduces the same rotation in the coordinates (q_0, q_{n+1}) .

We also remark the formal analogy between (6.5) and (A.3) in the Appendix.

Proof. Fix $w = (\zeta, v)$ and a tangent vector (η, u) at w . Set $b_\theta(w) = w_\theta = (\zeta_\theta, v_\theta)$. We compute

$$(b_\theta)_{*,w}(\eta, u) = \frac{d}{d\varepsilon}|_{\varepsilon=0} \left((c\mathbf{1} - s(\zeta + \varepsilon\eta))^{-1}(s\mathbf{1} + c(\zeta + \varepsilon\eta)), (c\mathbf{1} - s(\zeta + \varepsilon\eta))^{-1}(v + \varepsilon u) \right) ,$$

where $\varepsilon \in \mathbb{R}$, $c = \cos \theta$, $s = \sin \theta$, and prove that $\|(b_\theta)_{*,w}(\eta, u)\|_{w_\theta}^2$ does not depend on θ as long as $\zeta \neq (\cot \theta)\mathbf{1}$. It is convenient to restrict ourselves to $\zeta \notin \mathbb{R}\mathbf{1}$, $v \neq 0$, the general case following by continuity.

We take the first-order expansion of each component in ε . The expansion of the first component takes place in the subalgebra of (C, \cdot_{v_0}) (for some v_0) generated by ζ and η , which is associative. By standard computations, we find that

$$(6.6) \quad \begin{aligned} (b_\theta)_{*,w}(\eta, u) &\stackrel{\text{def}}{=} (\eta_\theta, u_\theta) \\ &= ((c\mathbf{1} - s\zeta)^{-1}\eta(c\mathbf{1} - s\zeta)^{-1}, (c\mathbf{1} - s\zeta)^{-1}u + s(c\mathbf{1} - s\zeta)^{-1}\eta(c\mathbf{1} - s\zeta)^{-1}v) . \end{aligned}$$

Notice that, according to Remark 2 in Section 1, the expression defining η_θ is independent of the choice of v_0 .

The subspace $C + Cw$ and its orthogonal complement $(C + Cw)^\perp$ (with respect to the product inner product on W) are clearly invariant under $(b_\theta)_{*,w}$. Notice that, since w_θ is in $C + Cw$, then $C + Cw$ is also the direct sum of Cw_θ and its orthogonal complement. By (6.1), $C + Cw$ and $(C + Cw)^\perp$ are also orthogonal with respect to the inner product $\langle \cdot, \cdot \rangle_{w_\theta}$. It is then sufficient to prove that $\|(\eta_\theta, u_\theta)\|_{w_\theta}^2$ does not depend on θ when (η, u) belongs to either of the two subspaces.

Take $(0, u) \in (C + Cw)^\perp$, i.e. with $u \perp Cv$ in V . Then $\eta_\theta = 0$ and $u_\theta = (-s\zeta + c\mathbf{1})^{-1}u$. Then

$$\|(0, u_\theta)\|_{w_\theta}^2 = \frac{|u_\theta|^2}{1 + |w_\theta|^2} = \frac{|u|^2}{(1 + |w_\theta|^2)|-s\zeta + c\mathbf{1}|^2} .$$

Observing that

$$(6.7) \quad 1 + |w_\theta|^2 = \frac{1 + |w|^2}{|c\mathbf{1} - s\zeta|^2} ,$$

the conclusion follows.

Take now $\xi = (\eta, \lambda\zeta^{-1}v) \in C + Cw$. Observe that $\xi \in Cw$ if and only if $\lambda = \eta$ and $\xi \perp Cw$ if and only if $\lambda = -\frac{|\zeta|^2}{|v|^2}\eta$. It follows that the orthogonal decomposition of ξ , for general η, λ , into a term in Cw and one orthogonal to it is

$$(\eta, \lambda\zeta^{-1}v) = (\alpha, \alpha\zeta^{-1}v) + \left(\beta, -\frac{|\zeta|^2}{|v|^2}\beta\zeta^{-1}v\right)$$

with

$$\begin{aligned}\alpha &= \frac{|\zeta|^2}{|w|^2}\eta + \frac{|v|^2}{|w|^2}\lambda, \\ \beta &= \frac{|v|^2}{|w|^2}(\eta - \lambda).\end{aligned}$$

Therefore, skipping the straightforward computations,

$$\begin{aligned}(6.8) \quad \|(\eta, \lambda\zeta^{-1}v)\|_{w+}^2 &= \frac{|\alpha|^2\left(1 + \frac{|v|^2}{|\zeta|^2}\right)}{(1 + |w|^2)^2} + \frac{|\beta|^2\left(1 + \frac{|\zeta|^2}{|v|^2}\right)}{1 + |w|^2} \\ &= \frac{1}{(1 + |w|^2)^2} (|\eta|^2 + |\lambda\zeta^{-1}v|^2 + |\eta - \lambda|^2|v|^2).\end{aligned}$$

Obviously, the same formula holds with w, v, ζ replaced by $w_\theta, v_\theta, \zeta_\theta$. We then apply (6.8) for a given θ , with $\eta = \eta_\theta$ as in (6.6) and $\lambda = \lambda_\theta$ such that $(\eta_\theta, \lambda_\theta\zeta_\theta^{-1}v_\theta) = (b_\theta)_{*,w}(\eta, \lambda\zeta^{-1}v)$, i.e.,

$$(6.9) \quad \lambda_\theta\zeta_\theta^{-1}v_\theta = (c\mathbf{1} - s\zeta)^{-1}(\lambda\zeta^{-1} + s\eta(c\mathbf{1} - s\zeta)^{-1})v.$$

By (6.7),

$$\frac{|\eta_\theta|^2}{(1 + |w_\theta|^2)^2} = \frac{|\eta|^2}{(1 + |w|^2)^2},$$

which does not depend on θ . The next term is

$$\frac{|\lambda_\theta\zeta_\theta^{-1}v_\theta|^2}{(1 + |w_\theta|^2)^2} = \frac{|c\mathbf{1} - s\zeta|^2|(\lambda\zeta^{-1} + s\eta(c\mathbf{1} - s\zeta)^{-1})v|^2}{(1 + |w|^2)^2}.$$

For the last term, we use the identity

$$|\eta_\theta - \lambda_\theta|^2|v_\theta|^2 = |\zeta_\theta|^2|\eta_\theta - \lambda_\theta|^2|\zeta_\theta^{-1}v_\theta|^2 = |\zeta_\theta|^2|(\eta_\theta - \lambda_\theta)\zeta_\theta^{-1}v_\theta|^2.$$

Simple calculations and the identity $(s\mathbf{1} + c\zeta)^{-1} - s\mathbf{1} = c(c\mathbf{1} - s\zeta)(s\mathbf{1} + c\zeta)^{-1}$ give that

$$(\eta_\theta - \lambda_\theta)\zeta_\theta^{-1}v_\theta = (c\mathbf{1} - s\zeta)^{-1}(c\eta(s\mathbf{1} + c\zeta)^{-1} - \lambda\zeta^{-1})v.$$

Hence

$$\frac{|\eta_\theta - \lambda_\theta|^2|v_\theta|^2}{(1 + |w_\theta|^2)^2} = \frac{|s\mathbf{1} + c\zeta|^2|(c\eta(s\mathbf{1} + c\zeta)^{-1} - \lambda\zeta^{-1})v|^2}{(1 + |w|^2)^2}.$$

We must then prove that

$$|c\mathbf{1} - s\zeta|^2 |(\lambda\zeta^{-1} + s\eta(c\mathbf{1} - s\zeta)^{-1})v|^2 + |s\mathbf{1} + c\zeta|^2 |(c\eta(s\mathbf{1} + c\zeta)^{-1} - \lambda\zeta^{-1})v|^2$$

does not depend on θ . Expanding the right-hand factor in each summand, this quantity equals

$$\begin{aligned} & (|c\mathbf{1} - s\zeta|^2 + |s\mathbf{1} + c\zeta|^2) |\lambda\zeta^{-1}v|^2 + (s^2 + c^2) |\eta|^2 |v|^2 \\ & \quad - 2|c\mathbf{1} - s\zeta|^2 s \langle \lambda\zeta^{-1}v, \eta(c\mathbf{1} - s\zeta)^{-1}v \rangle - 2|s\mathbf{1} + c\zeta|^2 c \langle \lambda\zeta^{-1}, v\eta(s\mathbf{1} + c\zeta)^{-1}v \rangle \\ & = (1 + |\zeta|^2) |\lambda\zeta^{-1}v|^2 + |\eta|^2 |v|^2 \\ & \quad + 2s \langle \lambda\zeta^{-1}v, \eta(c\mathbf{1} - s\bar{\zeta})v \rangle - 2c \langle \lambda\zeta^{-1}, v\eta(s\mathbf{1} + c\bar{\zeta})v \rangle \\ & = (1 + |\zeta|^2) |\lambda\zeta^{-1}v|^2 + |\eta|^2 |v|^2 - 2 \langle \lambda\zeta^{-1}v, \eta\bar{\zeta}v \rangle, \end{aligned}$$

a quantity that does not depend on θ . \square

Corollary 6.3. *The transition maps $\varphi_j \circ \varphi_k$ are isometries on $W \cap (\varphi_j \circ \varphi_k)^{-1}W$.*

Proof. Since $\varphi_0 = (-\text{id}) \circ b_{\frac{\pi}{2}}$, and $-\text{id} \in K$, φ_0 is an isometry on $W \setminus V$. Since $\varphi_j \in K$ for $1 \leq j \leq n$, then φ_j is an isometry on $W \setminus (Cv_j)^\perp$. Compositions are then isometries on appropriate open dense subsets of W and, by continuity, they remain isometric on $W \cap (\varphi_j \circ \varphi_k)^{-1}W$. \square

We can then extend the metric to CPW by imposing that the φ_j , $0 \leq j \leq n$, are isometries of the whole space. Observe that

- (i) by Proposition 6.1, W_∞ is the geodesic sphere centered at the origin of W and radius $\frac{\pi}{2}$;
- (ii) the b_θ extend uniquely to isometries of CPW ; for $\theta = 0, \pi$, $b_\theta \in K$, so that its extension is obvious; for $\theta \neq 0, \pi$ we have

$$\begin{aligned} & b_\theta((\cot \theta)\mathbf{1}, v) = [\mathbf{1}, (\sin \theta)v], \\ (6.10) \quad & b_\theta([\mathbf{1}, v]) = (-\cot \theta)\mathbf{1}, -(\sin \theta)^{-1}v, \\ & b_\theta([0, v]) = [0, v]; \end{aligned}$$

- (iii) $\gamma(\theta) = b_\theta(0)$ is the geodesic through 0 with tangent vector $(\mathbf{1}, 0)$ there, and its length is π .

Theorem 6.4. *CPW is a compact symmetric space of rank one.*

Proof. We show that CPW is homogeneous by proving that the origin can be mapped to any other point by isometries. If $w \in W$, take $\theta = \arctan|w|$, so that $b_\theta(0) = (|w|, 0)$. By Corollary 4.4, there is now $k \in K$ such that $k(|w|, 0) = w$. Take now $p = \pi(Cw) \in W_\infty$, and let $k \in K$ be such that $kC = Cw$. Then the extension \bar{k} of k to CPW maps $[\mathbf{1}, 0]$ into p . By (5.1), $\bar{k} \circ \varphi_0$ maps 0 into p .

The geodesic symmetry around 0 is $-\text{id}$, which is in K , hence CPW is symmetric. Since the action of K on the unit sphere in the tangent space at 0 is transitive, the rank of CPW is one. \square

Let U be the isometry group of CPW and $B = \{b_\theta : \theta \in \mathbb{T}\}$.

Corollary 6.5. $U = KBK$. The centralizer of B in K is M .

Proof. The proof of Theorem 6.4 shows that any isometry of CPW can be decomposed as $k_1 b_\theta k_2$ with $k_1, k_2 \in K$. Let $k \in K$ be such that $b_\theta k = k b_\theta$ for every θ . Then

$$((\tan \theta)\mathbf{1}, 0) = b_\theta k(0, 0) = k b_\theta(0, 0) = k((\tan \theta)\mathbf{1}, 0) ,$$

i.e. k fixes the points in $(\mathbf{1}, 0)$. By Proposition 4.1, $k \in M$. Conversely, take $m \in M$. Then $m = (\varphi, \psi)$, where φ, ψ satisfy (2.1). Given $(\zeta, v) \in W$, let $(\zeta', v') = (\varphi(\zeta), \psi(v))$. Set $c = \cos \theta, s = \sin \theta$. Hence

$$b_\theta(\zeta', v') = ((c\mathbf{1} - s\zeta')^{-1}(s\mathbf{1} + c\zeta'), (c\mathbf{1} - s\zeta')^{-1}v')$$

It follows from (2.1) that

$$\varphi(\eta \cdot_v \lambda) = \varphi(\eta) \cdot_{\psi(v)} \varphi(\lambda) .$$

By Remark 1 in Section 1, if η and λ are rational expression in ζ , the value of the product does not depend on v . We can then say that

$$(c\mathbf{1} - s\zeta')^{-1}(s\mathbf{1} + c\zeta') = \varphi((c\mathbf{1} - s\zeta)^{-1}(s\mathbf{1} + c\zeta)) ,$$

and

$$(c\mathbf{1} - s\zeta')^{-1}v' = \varphi((c\mathbf{1} - s\zeta)^{-1})\psi(v) = \psi((c\mathbf{1} - s\zeta)^{-1}v) ,$$

i.e., $b_\theta(\zeta', v') = m b_\theta(\zeta, v)$. \square

7. SOME APPLICATIONS

The purpose of this section is to use our setup to prove a few (known) fundamental facts about compact symmetric spaces of rank one. In the usual approach these things are easy to prove for spheres and classical projective spaces, but quite difficult for the octonionic case (cf. [Be], Ch. 3).

Throughout this section we consider a J^2C -module (C, V) and the associated $W = C \oplus V$, together with CPW . If V has a C -basis of n elements, we say that $\dim_C W = n + 1$.

Theorem 7.1. *If $d = \dim_{\mathbb{R}} C$ and $m = \dim_C W$, we have*

$$\text{vol}(CPW) = \frac{\Gamma(\frac{d}{2})}{\Gamma((m+1)\frac{d}{2})} \pi^{m\frac{d}{2}} .$$

Proof. Since W is open dense in CPW , by (6.1) we have

$$\text{vol}(CPW) = \int_W (1 + |w|^2)^{-2d - (m-1)d} dw ,$$

where dw is Lebesgue measure. Polar coordinates and the substitution $x = |w|^2$ reduce this to

$$\frac{1}{2} |S^{md-1}| \int_0^\infty (1+x)^{-(m+1)\frac{d}{2}} x^{m\frac{d}{2}-1} dx ,$$

which is a classical Beta-integral. \square

Theorem 7.2. *Let E be a plane in the tangent space to CPW at $(0,0)$ (identified with W as usual), and suppose that $X, Y \in W$ form an orthonormal basis of E . Then the sectional curvature $\sigma(E)$ of E is*

$$(7.1) \quad \sigma(E) = 1 + 3|\pi_{CX}Y|^2 = 1 + 3\cos^2\varphi ,$$

where π_{CX} denotes orthogonal projection onto CX and φ is the angle of CX and CY .

Proof. We use the classical formula

$$(7.2) \quad \sigma(E) = \lim_{r \rightarrow 0} \frac{3}{\pi} \frac{2\pi r - L_r}{r^3} ,$$

where L_r is the arc length of the exponential of a circle of radius r in E . By Proposition 6.1, this curve can be written as

$$\gamma_r(\theta) = \tan r((\cos \theta)X + (\sin \theta)Y) , \quad (0 \leq \theta \leq 2\pi) .$$

Since $\gamma_r(\theta)$ and $\dot{\gamma}_r(\theta) = \tan r(-(\sin \theta)X + (\cos \theta)Y)$ are images of $\gamma_r(0) = (\tan r)X$ and $\dot{\gamma}_r(0) = (\tan r)Y$ respectively under a real rotation, the angle φ of $C\gamma(\theta)$ and $C\dot{\gamma}_r(\theta)$ is independent of θ and $\cos \varphi = |\pi_{CX}Y|$ (cf. Remark 1 at the end of Section 4). Using the abbreviation $p = |\pi_{CX}Y|$ and (6.3), (6.4), it follows that

$$L_r = 2\pi \frac{\tan r}{1 + \tan^2 r} (1 + (1 - p^2) \tan^2 r)^{\frac{1}{2}} \sim 2\pi \left(r - \left(\frac{1}{6} + \frac{p^2}{2} \right) r^3 \right) ,$$

and (7.2) gives the result. \square

Remark. The C -lines of W have now the following geometric characterization: two linearly independent elements X, Y in W are in the same C -line through 0 if and only if $\sigma(E) = 4$ for the plane E spanned by them. Similarly, $CX \perp CY$ if and only if $\sigma(E) = 1$.

Next, we describe the totally geodesic submanifolds of CPW . We fix an o.n. C -basis $\{v_1, \dots, v_n\}$ of V and take a number n_0 , $0 \leq n_0 \leq n$. If $n_0 = 0$, let C_0 be any real subspace of C . If $n_0 \geq 1$, let C_0 be any division subalgebra of (C, \cdot_{v_1}) . In either case we set

$$(7.3) \quad W_0 = C_0 \oplus \sum_{j=1}^{n_0} C_0 v_j = C_0 \oplus V_0 .$$

We note that (C_0, V_0) is a J^2C -module. If $n > 1$, this follows from Proposition 1.4 and Corollary 1.5. If $n = 1$, it is trivial.

Theorem 7.3. *The closure of W_0 in CPW is totally geodesic. All totally geodesic submanifolds of CPW arise as U -images of these.*

Proof. It is well known and obvious that the fixed point set of an isometry in a Riemannian manifold is totally geodesic. The linear map which is the identity on $W_1 = C \oplus \sum_{j=1}^{n_0} C v_j$ and minus the identity on $\sum_{j=n_0+1}^n C v_j$ is in M , so it is an

isometry. Therefore W_1 is totally geodesic. It is then sufficient to prove that W_0 is totally geodesic in W_1 .

If $n_0 = 0$, there is not much to say. In this case $W_1 = C$. If C_0 is any subspace of C , the reflection with respect to C_0 is an isometry, and this implies that C_0 is totally geodesic. The closure of C in CPW is a d -dimensional sphere, and the closure of C_0 is a lower-dimensional geodesic sphere.

Assume therefore that $n_0 \geq 1$. Given a division subalgebra C_0 of C , there is a chain of division subalgebras between C_0 and C , each of index 2 in the next. So it is enough to consider the case where C_0 has index 2 in C . In this case we have the orthogonal direct sum $C = C_0 \oplus C_0z$, where z is an arbitrary unit element orthogonal to C_0 (the product being understood as \cdot_{v_1}).

Define $\alpha : C \rightarrow C$ by $\alpha(\zeta_1 + \zeta_2z) = \zeta_1 - \zeta_2z$, for $\zeta_1, \zeta_2 \in C_0$. Observe that left multiplication by a unit element $\eta \in C$ is orthogonal on C , which preserves the decomposition $C_0 \oplus C_0z$ if $\eta \in C_0$, and interchanges the two summands if $\eta \in C_0z$. Using this, it follows that α is an automorphism of C .

We extend α to W_1 by $\alpha(\zeta, \sum \zeta_j v_j) = (\alpha(\zeta), \sum \alpha(\zeta_j)v_j)$. This is an orthogonal transformation and it preserves C -lines through the origin. In fact, if $\zeta^{-1}\zeta_j v_j = \zeta'^{-1}\zeta'_j v_j$ for every j , then also $\alpha(\zeta)^{-1}\alpha(\zeta_j)v_j = \alpha(\zeta')^{-1}\alpha(\zeta'_j)v_j$, in the associative case obviously, and in the non-associative case because there is only one j .

Since W_0 is the fixed point set of α , this proves that W_0 is totally geodesic in W_1 , hence in W . Because the metric induced on W_0 from W is the same metric constructed starting from the J^2C -module (C_0, V_0) , the embedding of W_0 in CPW extends to an isometric embedding of C_0PW_0 , whose image is the closure of W_0 . This proves the first statement.

To prove the converse, let $N \subset CPW$ be totally geodesic. N is a symmetric space and clearly it has rank one. Translating by an element of U , we may assume that $0 \in N$. The tangent space to N at 0 is a subspace W_0 of W . Because geodesics through the origin are straight lines, N is the closure of W_0 in CPW .

By Theorem 8.4, whose proof is independent of the present theorem, there is a J^2C -module (C_0, V_0) such that N is isometric (up to a constant factor) to C_0PW_0 . Because C_0 has constant curvature equal to 4, it follows from the Remark above that $\dim C_0 \leq \dim C$. Acting by K , we can assume that $C_0 \subset C$. Similar considerations based on the same Remark imply that $V_0 \subset V$. Along the same lines, if $\{v_1, \dots, v_{n_0}\}$ is an orthonormal C_0 -basis of V_0 , the subspaces C_0v_j are contained in different, hence orthogonal, C -lines in V . Therefore $\{v_1, \dots, v_{n_0}\}$ can be completed to an orthonormal C -basis $\{v_1, \dots, v_n\}$ of V , and W_0 has the form (7.3). \square

We call *projective C -line* in CPW a maximal totally geodesic submanifold of constant curvature equal to 4. If no sectional curvature in CPW is equal to 4 (i.e. if $\dim C = 1$), we call projective C -lines the geodesics.

Corollary 7.4. *U acts transitively on projective C -lines. The projective C -lines intersecting W are the closures in CPW of affine C -lines in W . The projective C -lines contained in W_∞ are the images $\pi(E)$ of linear C -subspaces E of W of C -dimension two.*

Proof. The first statement follows immediately from Theorem 7.3.

Observe now that, for $0 \neq v \in V$, the affine C -line $(\tan \theta, 0) + Cv$ in W is the image of Cv under the isometry b_θ in (6.5). Therefore it is a geodesic in W , and the

same is true for its closure in CPW . By Corollary 6.5, this gives the description of all projective C -lines intersecting W .

One projective C -line entirely contained in W_∞ is obtained as the image under φ_0 of the projective C -line $Cv \cup \{[0, v]\}$ for a fixed $v \neq 0$. This gives $\{[1, \zeta v] : \zeta \in C\} \cup \{[0, v]\}$, equal to $\pi(C \oplus Cv)$. If we act on this line by an element $u = k_1 b_\theta k_2$ of U , the image will remain inside W_∞ if and only if $b_\theta \pi(k_2(C \oplus Cv)) \subset W_\infty$. By (6.10), if E is a C -subspace of dimension 2 and b_θ maps $\pi(E)$ into W_∞ , then necessarily $\theta \in \{0, \pi\}$, so that $u \in K$.

It follows that all the projective C -lines in W_∞ are the images of $C \oplus Cv$ under the action of K , and the conclusion follows from Corollary 4.7. \square

It is clear that any two distinct points of CPW determine a unique projective C -line. We also note that the projective C -lines are the ‘‘Helgason spheres’’ which, by [Hel2], are present in any compact symmetric space.

Generalizing the notion of projective C -line, we call *projective C -subspace* a subset E of CPW such that for any pair of distinct points of E , the whole projective C -line joining them is contained in E . If a projective C -subspace E intersects W , it is easy to see that $E \cap W$ is an affine C -subspace of W . Since we can always put ourselves in this situation applying an element of U , it follows that E is a submanifold of real dimension equal to a multiple kd of d . We then say that E has C -dimension equal to k . By Theorem 7.3, the projective C -subspaces are the U -images of $W_0 \cup \pi(W_0)$, with W_0 as in (7.3) and $C_0 = C$. We also have the following analogue of Corollary 7.4.

Corollary 7.5. *U acts transitively on k -dimensional projective C -subspaces. The k -dimensional projective C -subspaces intersecting W are the closures in CPW of affine C -subspaces in W . The k -dimensional projective C -subspaces contained in W_∞ are the images $\pi(E)$ of linear C -subspaces E of W of C -dimension $k + 1$.*

A number of further geometric facts follow very easily. For instance, the Jacobi fields along a geodesic can be determined by reduction to the geodesic $\gamma(t) = (\tan t, 0)$. For $z \in C'$ and $v \in V$, let $z(t)$ and $v(t)$ be their parallel displacements along γ . Because C is totally geodesic in W , $z(t)$ is the same as parallel displacement in C of z along γ . Since C has constant curvature 4, it follows (cf. [KN, vol. I, p. 71]) that $(\sin 2t)z(t)$ and $(\cos 2t)z(t)$ are Jacobi fields. Similarly, $v(t)$ is the same as parallel displacement in $\mathbb{R}\mathbf{1} \oplus \mathbb{R}v$ of v along γ . Therefore $(\sin t)v(t)$ and $(\cos t)v(t)$ are Jacobi fields. By a dimension count, all Jacobi fields along γ are linear combination of these and of $\dot{\gamma}(t)$ and $t\dot{\gamma}(t)$.

Another fact concerns cut locus and conjugate points of a given point $p \in CPW$. We can assume that $p = 0$. As we have already observed, two geodesics, $\exp_0(tX)$ and $\exp_0(tY)$, meet only at 0 if X and Y are not in the same C -line, and at the two points 0 and $[X] = [Y]$ if X and Y belong to the same C -line. It is then clear that the cut locus of 0 is W_∞ , or, in other words, the set of points q such that $d(0, q) = \frac{\pi}{2}$. In the latter formulation, the statement is true for any p in place of 0.

It also follows that the conjugate locus of any point p is $\{p\}$ when $d = \dim C = 1$ (the real projective space), and coincides with the cut locus otherwise. In the latter case the multiplicity of the first conjugate point along any geodesic is $d - 1$.

Finally we note that CPW has a natural cell decomposition

$$CPW = \mathcal{C}^{d(n+1)} \cup \mathcal{C}^{dn} \cup \dots \cup \mathcal{C}^0,$$

where \mathcal{C}^k denotes a cell of dimension k . In fact, we can take $\mathcal{C}^{d(n+1)} = W$. Its complement W_∞ is isometric, under either of the maps φ_j in (5.2), with the closure of $C \oplus (Cv_j)^\perp = W_j$. This closure is CPW_j , with one less C -dimension. So the statement follows by induction.

8. EVERY COMPACT RANK-ONE SYMMETRIC SPACE IS A CPW

The reader who is willing to accept the classification list of symmetric spaces existing in the literature, e.g. [Be, Hel1, W], can compare that list with the list in Section 3 of J^2C -modules and the corresponding CPW 's, and convince himself that we have obtained all compact rank-one symmetric spaces. However, in this section we will give a classification-free proof of this fact, clarifying at the same time the duality relations between compact and non-compact spaces.

Lemma 8.1. *CPW is simply connected if and only if $d = \dim C > 1$.*

Proof. If $d = 1$, then CPW is $\mathbb{R}P^{n+1}$ if $n = \dim V > 1$ and S^1 if $V = 0$. Therefore CPW is not simply connected.

Suppose now that $d > 1$, and observe that if $\dim W > 1$ and W_∞ is simply connected, so is CPW . This depends on the fact that W_∞ is a deformation retract of CPW minus a point in W .

If V is non-trivial and v_1 is a unit vector in V , set $V' = (Cv_1)^\perp$. The map φ_1 in (5.2) establishes a diffeomorphism between W_∞ and CPW' (cf. end of Section 7), with $W' = C \oplus V'$. By induction, matters are reduced to $V = 0$. In this case CPW is the sphere S^d , which is simply connected. \square

It is well known (cf. [W]) that the duality between compact and non-compact orthogonal semisimple Lie algebras induces a (bijective) duality between simply connected compact symmetric spaces and symmetric spaces of the non-compact type. Duality respects the rank, and the balls B_W described in the Appendix, with the metric defined in (A.1), give us models of all the rank-one symmetric spaces of non-compact type. In fact, we can restrict ourselves to those $W = C \oplus V$ with $d = \dim C > 1$.

Proposition 8.2. *Let $W = C \oplus V$ with $d > 1$. The simply connected compact dual of the unit ball B_W is CPW .*

Proof. Assume that $d > 1$. The non-compact dual of CPW must be a ball B_{W^\sharp} for some $W^\sharp = C^\sharp \oplus V^\sharp$ with the same dimension as W . Take the origin in W as base points in CPW and the origin in W^\sharp as base point in B_{W^\sharp} .

By [W, Cor. 8.4.3], the two tangent spaces can be identified in such a way that sectional curvatures of corresponding planes have opposite values, possibly up to a constant factor. It follows from Theorem 7.2 and the comments following Lemma A.1 in the Appendix that C and C^\sharp must have the same dimension. Then $W^\sharp = W$. \square

It follows that the CPW with $d > 1$ are all the simply connected compact symmetric spaces of rank one. Setting aside the trivial one-dimensional case ($C =$

\mathbb{R} , $V = 0$), every other compact symmetric spaces of rank one must have a *CPW* with $d > 1$ as its simply connected covering. We are so led to discuss the existence of locally isometric symmetric quotients of the *CPW*.

Lemma 8.3. *Suppose that CPW is a non-trivial covering of a symmetric space X . Then $V = 0$ and CPW is a two-fold covering of X .*

Proof. Let $\pi : CPW \rightarrow X$ be the covering map. Since X has rank one, its geodesics are circles and they all have the same length. Let E be the set of points $p \in CPW$, different from 0, that are mapped to $\pi(0)$. Then E is non-empty and finite. Take $p_0 \in E$ of minimal distance from 0 and let γ be a full geodesic circle in *CPW* going through 0 and p . Then $\pi(\gamma)$ is a geodesic circle in X and its length is $\delta = d(0, p_0)$, strictly smaller than the length of γ . It follows that every geodesic circle in *CPW* containing 0 is mapped by π onto a geodesic circle of length δ , hence it must contain a point in E at distance δ from 0.

Since E is finite, we surely have two distinct geodesics through 0 intersecting at a point $p \in E$. By Proposition 5.2, geodesics through zero have the form $\mathbb{R}w \cup [w]$ with $w \in W$. Therefore two distinct geodesics can have a common point different from 0 only if they lie in the same C -line Cw , and in this case the common point is $[w]$. This implies that $\delta = \frac{\pi}{2}$ and hence $[w] \in E$ for every $w \in W$. So W_∞ is finite. But this is only possible if $W = C$, and then $E = W_\infty$ consists of one single point. \square

Theorem 8.4. *The CPW are all the distinct compact symmetric spaces of rank one, including the circle S^1 .*

Proof. It remains to verify that if $d = 1$ and $W = \mathbb{R} \oplus V$ with $V \neq 0$, then *CPW* is isometric, up to a factor, to the quotient of $CP\tilde{W}$ modulo the antipodal map, with $\tilde{C} = W$ and $\tilde{V} = 0$. To see this, consider the map π from \tilde{W} to W given by

$$\pi(w) = \frac{2w}{1 - |w|^2}, \quad (|w| \neq 1).$$

This is an isometry up to a factor 2, and it extends continuously to the unit sphere and to the point at infinity, identifying the pairs of antipodal points, w and $-\frac{w}{|w|^2}$. \square

9. THE GROUP $GL(W, C)$

We study $GL(C, W)$ in some detail and, to avoid trivialities, we assume that $V \neq 0$.

$GL(W, C)$ is closed under adjoints. To see this, note that for any \mathbb{R} -subspace W_1 , $w \perp gW_1$ if and only if $g^*w \perp W_1$. Hence $g^*((gW_1)^\perp) = W_1^\perp$. Now let Cw be a C -line and let $W_1 = g^{-1}((Cw)^\perp)$. By Lemma 4.5, $(Cw)^\perp$ is a C -subspace, and by Proposition 4.6 so is W_1 . By Lemma 4.5 again, $g^*(Cw)$ is a C -line.

It follows that $GL(W, C)$ is a reductive Lie group, since it is fixed under the Cartan involution $g \mapsto g^{*-1}$ of $GL(W, \mathbb{R})$ (cf. [W]). What follows is a useful characterization of the elements of $GL(W, C)$.

Theorem 9.1. *Assume that $V \neq 0$. The elements of $GL(W, C)$ are the homeomorphisms of W onto itself fixing the origin and mapping parallel C -lines into parallel C -lines.*

Proof. One implication is obvious. So let g be a homeomorphism fixing 0 and mapping parallel C -lines into parallel C -lines. We need to prove that g is \mathbb{R} -linear. Since g is continuous, it suffices to prove that it is additive.

If w, w' are points in W belonging to different C -lines through 0, it is easily verified that $w + w'$ is the only point in the intersection of $w + Cw'$ with $w' + Cw$.

Let w, w' be as above. By assumption, $g(Cw) = Cg(w)$ and $g(w' + Cw) = g(w') + Cg(w)$. Similarly, $g(Cw') = Cg(w')$ and $g(w + Cw') = g(w) + Cg(w')$. By injectivity, $Cg(w)$ and $Cg(w')$ are different C -lines. Therefore $g(w + w') \in (g(w') + Cg(w)) \cap (g(w) + Cg(w'))$. By the previous remark, $g(w + w') = g(w) + g(w')$.

If w, w' belong to the same C -line through 0, the same identity follows by continuity, using the existence of other C -lines. \square

By Corollary 4.7, every $g \in GL(W, C)$ can be decomposed as $g = k \circ h$ with $k \in K$ and $h \in GL(W, C)$ such that $h(V) = V$.

Lemma 9.2. *Assume that $h \in GL(W, C)$ maps V into itself and let*

$$(9.1) \quad h = \begin{pmatrix} \alpha & 0 \\ \alpha(\cdot)v_0 & \varphi \end{pmatrix},$$

be the matrix representing it relative to the decomposition $W = C \oplus V$. Then

- (i) $\alpha : C \rightarrow C$ is a scalar multiple of an orthogonal transformation,
- (ii) $\varphi : V \rightarrow V$ is invertible and satisfies $\varphi(\zeta v) = \alpha(\zeta)\alpha(\mathbf{1})^{-1}\varphi(v)$,
- (iii) $v_0 \in V$.

Conversely, every triple (α, φ, v_0) satisfying (i), (ii), (iii) defines, through (9.1), an element of $GL(W, C)$ mapping V into itself.

Proof. Any h mapping V into itself is represented by a matrix

$$h = \begin{pmatrix} \alpha & 0 \\ \sigma & \varphi \end{pmatrix}.$$

The condition $h(C)$ equal to a C -line implies that

$$\alpha(\zeta)^{-1}\sigma(\zeta) = \alpha(\mathbf{1})^{-1}\sigma(\mathbf{1}) = v_0,$$

i.e. $\sigma(\zeta) = \alpha(\zeta)v_0$.

For $v \in V$, impose now that $h(0, \zeta v) \sim h(0, v)$. This gives $\varphi(\zeta v) = \beta(\zeta, v)\varphi(v)$, with $\beta(\zeta, v) \in C$ and $\beta(\mathbf{1}, v) = 1$.

Imposing now that

$$\psi(\zeta, \zeta v) = (\alpha(\zeta), \alpha(\zeta)v_0 + \beta(\zeta, v)\varphi(v)) \sim \psi(\mathbf{1}, v) = (\alpha(\mathbf{1}), \alpha(\mathbf{1})v_0 + \varphi(v)),$$

we obtain that

$$v_0 + \alpha(\zeta)^{-1}\beta(\zeta, v)\varphi(v) = v_0 + \alpha(\mathbf{1})^{-1}\varphi(v),$$

i.e. $\beta(\zeta, v) = \alpha(\zeta)\alpha(\mathbf{1})^{-1}$.

If $z \in C'$ and $|z| = 1$, then $z^2 = -\mathbf{1}$, and

$$\varphi(v) = -\varphi(z^2v) = -(\alpha(z)\alpha(\mathbf{1})^{-1})^2\varphi(v).$$

We then have $|\alpha(z)| = |\alpha(\mathbf{1})|$ and $\alpha(z) \cdot_{\varphi(v)} \alpha(\mathbf{1})^{-1} \in C'$. In particular $\alpha(z) \perp \alpha(\mathbf{1})$ for $z \in C'$. It follows that $|\alpha(\zeta)| = |\alpha(\mathbf{1})|$ for every $\zeta \in C$.

To proof of the converse is easy and we leave it to the reader. \square

An immediate consequence of this lemma is the following statement, which will be used in Section 10.

Corollary 9.3. *Let $g \in GL(W, C)$. The restriction of g to any C -line Cw through the origin is a scalar multiple of an orthogonal map.*

Proof. In order to prove this, decompose $g = kh$ with $k \in K$ and h as in Lemma 9.2. It is then sufficient to assume that $g = h$. By density, we can also assume that $w = (\mathbf{1}, v)$. Take $w' = (\zeta, \zeta v) \in Cw$. Then $|w'| = |\zeta||w|$ and

$$\begin{aligned} |h(w')|^2 &= |\alpha(\zeta)|^2 + |\alpha(\zeta)v_0 + \alpha(\zeta)\alpha(\mathbf{1})^{-1}\varphi(v)|^2 \\ &= |\alpha(\zeta)|^2|\alpha(\mathbf{1})|^{-2}(|\alpha(\mathbf{1})|^2 + |\alpha(\mathbf{1})v_0 + \varphi(v)|^2)^2 \\ &= |\zeta|^2|h(w)|^2. \end{aligned}$$

Hence the quantity $|h(w')|/|w'|$ is constant on Cw . \square

We fix now an ordered orthonormal C -basis $\{w_0, \dots, w_n\}$ of W with $w_0 = (\mathbf{1}, 0)$ and the other $w_j = (0, u_j)$ in V . Then any element w of W can be written as

$$w = \sum_{j=0}^n \zeta_j w_j,$$

with the same abuse of notation as in Proposition 4.2. This basis induces a flag $\{W_j\}$ of linear C -subspaces in W , with

$$(9.2) \quad W_j = \text{span} \{w_j, \dots, w_n\}.$$

Let P be the subgroup of $GL(W, C)$ consisting of the elements h that preserve the flag, i.e. such that $h(W_j) = W_j$ for every j .

We introduce three subgroups of P . The first group is $M_P = P \cap K = P \cap L$. By Corollary 4.7, M_P acts transitively on the product of unit spheres in the various Cw_j . The second group $\mathcal{A} \cong \mathbb{R}^{n+1}$ acts as scalar multiplication by $t_j > 0$ on each Cw_j . The third is the group N of those $h \in P$ such that $h|_{Cw_j} = \text{id} \pmod{W_{j+1}}$ for each j .

It is easy to verify that the elements of N are in one-to-one correspondence with the lower-triangular matrices with entries $\lambda_{ij} \in C$,

$$(9.3) \quad \Lambda = \begin{pmatrix} 1 & 0 & \dots & \dots & 0 \\ \lambda_{10} & 1 & 0 & \dots & 0 \\ \lambda_{20} & \lambda_{21} & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ \lambda_{n0} & \lambda_{n1} & \dots & \dots & 1 \end{pmatrix},$$

in the sense that to each Λ as above we associate $n \in N$ given by

$$n \left(\sum_{j=0}^n \zeta_j w_j \right) = \sum_{j=0}^n \left(\sum_{k < j} \zeta_k \cdot_{v_j} \lambda_{jk} + \zeta_j \right) w_j .$$

The product $n_1 n_2$ in N corresponds to the matrix product ${}^t(\Lambda_2 {}^t\Lambda_1)$ (the double transposition is a consequence of the fact that we write the action of C on V as a left action). Notice that, if C is associative, multiplication in C is unambiguously defined, whereas, if C is non-associative, the matrices Λ are 2×2 and their product only involves the sum of their $(1, 0)$ -entries.

We then have the following Langlands decomposition of P and Iwasawa decomposition of $GL(W, C)$.

Proposition 9.4. *P is the semidirect product $M_P \mathcal{A} N$. M_P and \mathcal{A} commute, and $M_P \mathcal{A}$ normalizes N . Moreover, $GL(W, C) = K \mathcal{A} N$.*

Proof. It is clear that M_P and \mathcal{A} commute, and that $M_P \mathcal{A}$ normalizes N .

Given $h \in P$, let α, φ, v_0 be as in Lemma 9.2.

For $j \geq 1$, $h(w_j) = (0, \varphi(u_j)) \in W_j$. Invertibility of φ implies that the (Cu_j) -component of $\varphi(u_j)$ must be different from zero. We can then set

$$(9.4) \quad \varphi(u_j) = \eta_j (u_j + \lambda_{j+1,j} u_{j+1} + \cdots + \lambda_{nj} u_n) \quad (\forall j \geq 1) ,$$

with $\eta_j \neq 0$. If

$$(9.5) \quad \begin{aligned} w'_0 &= (\mathbf{1}, v_0) , \\ w'_j &= (0, u_j + \lambda_{j+1,j} u_{j+1} + \cdots + \lambda_{nj} u_n) \quad (\forall j \geq 1) , \end{aligned}$$

then h maps each C -line Cw_j onto Cw'_j . Let

$$(9.6) \quad v_0 = \lambda_{10} u_1 + \cdots + \lambda_{n0} u_n ,$$

and let Λ be the lower-triangular matrix with all 1's along the diagonal and the λ_{ij} defined in (9.4) and (9.6). The corresponding element $n \in N$ also maps each C -line Cw_j onto Cw'_j . Therefore $n^{-1}h$ maps Cw_j into itself for every j .

Precisely, one verifies that

$$\begin{aligned} n^{-1}h(\zeta, 0) &= (\alpha(\zeta), 0) , \\ n^{-1}h(0, \zeta u_j) &= (0, \alpha(\zeta) \alpha(1)^{-1} \eta_j u_j) \quad (\forall j \geq 1) . \end{aligned}$$

It follows that $n^{-1}h$ is a scalar multiple of an orthogonal transformation on each C -line Cw_j , hence $n^{-1}h = ma \in M_P \mathcal{A}$. So $h = nma$, and, since $M_P \mathcal{A}$ normalizes N , $h = man'$ for some other $n' \in N$.

To show uniqueness, it suffices to observe that $M_P \cap \mathcal{A}$ and $(M_P \mathcal{A}) \cap N$ are trivial.

We have already pointed out that any element of $GL(W, C)$ is the product of an element of K and an element of P . Therefore $g \in GL(W, C)$ can be decomposed as kan , with $k \in K$, $a \in \mathcal{A}$, $n \in N$. If $kan = k'a'n'$, then $k^{-1}k' = an(a'n')^{-1}$ is the identity on each Cw_j , hence $k = k'$. Then $a = a'$ and $n = n'$, by the previous part of the proof. \square

Finally, we easily obtain the Cartan decomposition of $GL(W, C)$.

Theorem 9.5. $GL(W, C) = KAK$.

Proof. Given $g \in GL(W, C)$, g^*g is C -linear, self-adjoint and positive. Its eigenspaces are C -subspaces, so there is an orthonormal C -basis $\{e_0, \dots, e_n\}$ of W such that $g^*g = t_j^2 \text{id}$ on Ce_j for every j . Take $k \in K$ such that $ke_j = w_j$, with w_j as above, and let $a \in \mathcal{A}$ be such that $a|_{Cw_j} = t_j \text{id}$ for every j . Then $a^{-1}kg^*gk^{-1}a^{-1} = \text{id}$, i.e. $gk^{-1}a^{-1} \in K$. \square

10. THE GROUP OF COLLINEATIONS

We consider here a larger group of transformations of CPW than the isometry group U , i.e. the group of homeomorphisms that preserve the class of projective C -lines. These transformations are called *collineations* in projective geometry. Clearly, this definition makes sense only if CPW is not reduced to one single C -line, i.e. if $V \neq 0$. As we will see in Corollary 10.4, the restriction of a collineation to a single C -line turns out to be a conformal map. For this reason, when $V = 0$ the *conformal group* is the natural substitute of the collineation group. In the following we will assume that V is non-trivial.

The elements of U are obviously collineations and such are translations by Corollary 7.4. We remark in the Appendix that the group of collineations also includes the isometry group of the non-compact dual B_W of CPW . Collineations obviously map projective C -subspaces into projective C -subspaces of the same dimension.

Denote by \mathcal{G} the collineation group. If g is a collineation, there exist $u \in U$ such that $u(W_\infty) = g(W_\infty)$. Then there is a translation τ such that

$$(10.1) \quad g = u\tilde{g}\tau,$$

with \tilde{g} a collineation fixing 0 and stabilizing W_∞ . It follows from Theorem 9.1 that \tilde{g} is the continuous extension of an element of $GL(W, C)$. More generally, we can say that the collineations stabilizing W_∞ are the extensions of C -affine invertible transformations of W .

It follows from (10.1) that collineations map projective C -subspaces into projective C -subspaces of the same dimension.

Let $\{W_j\}$ be the flag (9.2) of linear C -subspaces of W , and consider the chain of projective C -subspaces of CPW

$$\pi(W_n) \subset \dots \subset \pi(W_0) = W_\infty.$$

Let $P_{\mathcal{G}}$ be the subgroup of \mathcal{G} whose elements preserve each $\pi(W_j)$. Then (up to extension to CPW) $P_{\mathcal{G}}$ is the semidirect product of P , introduced in Section 5, with the translation group $\mathcal{W} \cong W$. Let $N_{\mathcal{G}}$ be the nilpotent group given by the semidirect product of $N \subset P$ with \mathcal{W} . We then have the following decomposition of \mathcal{G} .

Theorem 10.1. *Any $g \in \mathcal{G}$ can be uniquely decomposed as $g = uan$, with $u \in U$, $a \in \mathcal{A}$ and $n \in N_{\mathcal{G}}$.*

Proof. Let $g = u\tilde{g}\tau$ as in (10.1). By Proposition 9.4, $\tilde{g} = kan_0$ with $k \in K$, $a \in \mathcal{A}$, $n_0 \in N$. Then $g = (uk)a(n_0\tau) \in U\mathcal{A}N_{\mathcal{G}}$.

If now $uan = u'a'n'$, with $u, u' \in U$, $a, a' \in \mathcal{A}$, $n, n' \in N_{\mathcal{G}}$, then $u^{-1}u' = an(a'n')^{-1} \in P_{\mathcal{G}}$. Writing $n = n_0\tau$, $n' = n'_0\tau'$, with $n_0, n'_0 \in N$, $\tau, \tau' \in \mathcal{W}$, the fact that $an(a'n')^{-1} \in GL(W, C)$ implies that $\tau = \tau'$. Therefore $u^{-1}u' = an_0(a'n'_0)^{-1} \in P$. Hence $u^{-1}u' \in K \cap P$ and $u = u'$. Finally, $an_0 = a'n'_0$ implies that $a = a'$, $n_0 = n'_0$. \square

We introduce now an involution on \mathcal{G} . As observed in Section 7, W_{∞} is the set of points at distance $\frac{\pi}{2}$ from the origin of W , also characterized as the cut locus of 0. By U -invariance, it follows that the cut locus p^* of any point $p \in CPW$ is a projective C -hyperplane, and that every projective C -hyperplane is the cut locus of some point. The correspondence $p \mapsto p^*$ is a bijection between points and projective C -hyperplanes.

Lemma 10.2. *The correspondence $p \mapsto p^*$ is such that*

$$(10.2) \quad \begin{aligned} 0^* &= W_{\infty}, \\ w^* \cap W &= (Cw)^{\perp} - \frac{1}{|w|^2}w && (\forall w \in W \setminus \{0\}), \\ [w]^* \cap W &= (Cw)^{\perp} && (\forall [w] \in W_{\infty}). \end{aligned}$$

To each collineation g we can associate the map $\theta g : CPW \rightarrow CPW$ defined by

$$(\theta g(p))^* = g(p^*).$$

Then

- (i) $\theta g = (g^*)^{-1}$ if $g \in GL(W, C)$,
- (ii) for every g , θg is a collineation,
- (iii) θ is an involutive automorphism of \mathcal{G} ,
- (iv) $\theta g = g$ if and only if $g \in U$.

Proof. For $g \in U$, we trivially have that $g(p^*) = (g(p))^*$ (i.e. $\theta g = g$). Modulo the action of K , it is then sufficient to identify $(t\mathbf{1}, 0)^*$ for $t \in \mathbb{R}$ and $[\mathbf{1}, 0]^*$ in order to obtain (10.2). Let b_s , $s \in \mathbb{T}$, be the maps defined in (6.5). Since $b_s(0) = (\tan s \mathbf{1}, 0)$ for $s \neq \pm \frac{\pi}{2}$ and $b_{\pm \frac{\pi}{2}}(0) = [\mathbf{1}, 0]$, the problem reduces to determining $b_s(W_{\infty})$. By (6.10),

$$b_s(W_{\infty}^1) = \left\{ \left(-(\cot s)\mathbf{1}, -\frac{1}{\sin s}v \right) : v \in V \right\} = -((\cot s)\mathbf{1}) + V,$$

and $(t\mathbf{1}, 0)^*$ is its closure. Taking $s = \frac{\pi}{2}$, this gives $[\mathbf{1}, 0]^*$.

The map θg is well defined for $g \in \mathcal{G}$. If $g \in U$, $g(p^*) = (g(p))^*$, i.e. $\theta g = g$. Moreover, for every $g, h \in \mathcal{G}$,

$$((\theta g)(\theta h)(p))^* = g((\theta h(p))^*) = gh(p^*) = (\theta(gh)(p))^*,$$

i.e. $(\theta g)(\theta h) = \theta(gh)$.

To prove (i), fix $w \in W$, $w \neq 0$, and take $k \in K$ such that $kw = (t\mathbf{1}, 0)$ with $t > 0$. Writing $g = g'k$, we have $\theta g(w) = (\theta g')kw = \theta g'(t\mathbf{1}, 0)$. Decompose g' as $k'h$ with $k' \in K$ and h as in (9.1). Then $\theta g'(t\mathbf{1}, 0) = k'\theta h(t\mathbf{1}, 0)$.

By (10.2),

$$h((t\mathbf{1}, 0)^*) = h(V - (t^{-1}\mathbf{1}, 0)) = V - (t^{-1}\alpha(\mathbf{1}), 0),$$

i.e. $\theta h(t\mathbf{1}, 0) = (t\alpha(\mathbf{1})|\alpha(\mathbf{1})|^{-2}, 0) = (h^*)^{-1}(t\mathbf{1}, 0)$. Finally,

$$\theta g(w) = \theta(k' h k)(w) = k'(\theta h)k(w) = k'(h^*)^{-1}k(w) = (g^*)^{-1}(w) .$$

To prove (ii), by (10.1) it suffices to prove that, for any translation $\tau_{w_0}(w) = w + w_0$, $\theta\tau_{w_0}$ is a collineation. Since for $w \in W$, $w \neq 0$,

$$\tau_{w_0}(w^*) = (Cw)^\perp - \frac{1}{|w|^2}w + w_0 = (Cw)^\perp - \frac{1}{|w|^2}w + P_{Cw}w_0 ,$$

with P denoting orthogonal projection, $\theta\tau_{w_0}(w)$ is the element $w' \in Cw$ such that

$$\frac{1}{|w'|^2}w' = \frac{1}{|w|^2}w - P_{Cw}w_0 .$$

Take $k \in K$ such that $kw_0 = (t\mathbf{1}, 0)$ with $t > 0$. Then $\tau_{w_0} = k^{-1}\tau_{kw_0}k$ and $\theta\tau_{w_0} = k^{-1}(\theta\tau_{kw_0})kw$. We can then restrict ourselves to $w_0 = (t\mathbf{1}, 0)$.

We compute $P_{Cw}(\mathbf{1}, 0)$ for $w = (\eta, u)$, assuming that $\eta \neq 0$ (for $\eta = 0$ the projection is 0, and $w' = w$). We impose that

$$(\mathbf{1} - \lambda, -\lambda\eta^{-1}u) \perp (\alpha, \alpha\eta^{-1}u) \quad (\forall \alpha \in C)$$

and obtain that

$$P_{Cw}(\mathbf{1}, 0) = \frac{1}{|w|^2}(|\eta|^2, \bar{\eta}u) .$$

With standard simplifications, this gives

$$(10.3) \quad \theta\tau_{w_0}(\eta, u) = ((\mathbf{1} - t\eta)^{-1}\eta, (\mathbf{1} - t\eta)^{-1}u) .$$

To prove that this is a collineation, we observe that (10.3) admits a continuous bijective extension to CPW . By continuity, we can limit ourselves to verify that the image of an affine C -line in W that is not parallel to V is contained in a projective C -line. Therefore we take

$$w = \omega_\zeta = (\zeta_0 + \zeta, v_0 + \zeta v_1) ,$$

with ζ varying in C and ζ_0, v_0, v_1 fixed. We show that the points

$$\theta\tau_{w_0}(\omega_\zeta) - \theta\tau_{w_0}(\omega_0) = (\lambda_\zeta, u_\zeta) ,$$

with

$$\begin{aligned} \lambda_\zeta &= (\mathbf{1} - t(\zeta_0 + \zeta))^{-1}(\zeta_0 + \zeta) - (\mathbf{1} - t\zeta_0)^{-1}\zeta_0 \\ u_\zeta &= (\mathbf{1} - t(\zeta_0 + \zeta))^{-1}(v_0 + \zeta v_1) - (\mathbf{1} - t\zeta_0)^{-1}v_0 \end{aligned}$$

lie on the same C -line through 0. A straightforward computation shows that $\lambda_\zeta^{-1}u_\zeta$ is independent of ζ .

Concerning (iii), we have already observed in the course of this proof that θ is multiplicative. By (10.1), (i) and (ii), θ maps \mathcal{G} into itself. The identity $\theta^2 g = g$ holds for $g \in U$ obviously and for $g \in GL(W, C)$ by (i). Computing $\theta^2\tau_{w_0}$ from (10.3), we conclude that it also holds for translations.

Finally, assume that $\theta g = g$. Modulo U , we can assume that $g(0) = 0$. Therefore, $g(W_\infty) = (\theta g(0)^*) = W_\infty$. Therefore $g \in GL(W, C)$ and $g = (g^*)^{-1}$, i.e. $g \in K$. \square

Theorem 10.3. \mathcal{G} is a simple Lie group G with restricted root system of type A_{n+1} and the differential of θ (also denoted by θ) is a Cartan involution on its Lie algebra. The corresponding Cartan decomposition of the Lie algebra \mathfrak{g} of \mathcal{G} is $\mathfrak{g} = \mathfrak{u} + \mathfrak{p}$, with \mathfrak{u} the Lie algebra of U . The Lie algebra \mathfrak{a} of \mathcal{A} is maximal abelian in \mathfrak{p} and the decomposition in Theorem 10.1 is the associated Iwasawa decomposition.

Proof. We sketch two proofs of the fact that \mathcal{G} is a Lie group. The first one makes use of the theorem of Montgomery-Zippin-Gleason. Since \mathcal{G} with the topology of uniform convergence on CPW is clearly a topological transformation group, it suffices to prove that \mathcal{G} is locally Euclidean.

The stabilizer of 0 in \mathcal{G} of W_∞ as a set is the semidirect product $GL(W, C) \times_s \mathcal{W}$. Hence the stabilizer of 0 is its θ -image $B = GL(W, C) \times_s \theta\mathcal{W}$. Since θ is easily seen to be a homeomorphism of \mathcal{G} , B is locally Euclidean with the topology induced from \mathcal{G} . Since $\mathcal{W} \cdot 0 = W$, it follows that $\mathcal{W}B$ is a neighborhood of the identity in \mathcal{G} . If $U_{\mathcal{W}}$ and U_B are small neighborhoods of the identity in \mathcal{W} and B respectively, then $U_{\mathcal{W}}U_B$ is a neighborhood of the identity in \mathcal{G} and is homeomorphic to $U_{\mathcal{W}} \times U_B$, showing that \mathcal{G} is locally Euclidean.

The second proof consists in introducing local coordinates near the identity coming from the decomposition $\mathcal{W} \cdot GL(W, C) \cdot \theta\mathcal{W}$ of an open dense subset of \mathcal{G} . The map $g \mapsto \theta g^{-1}$ is clearly smooth on this set, and one needs to verify smoothness of the group operations. The key point is then to verify, with the aid of the formulas defining θ , that the map $(w_0, w) \mapsto \theta\tau_{w_0}(w)$ of $W \times W$ into W is rational, and therefore smooth where meaningful. From this one gets that θ and the inversion are both smooth on a neighborhood of the identity. With appropriate use of left translations, one finishes the proof.

The existence of the Cartan involution θ implies that \mathfrak{g} is reductive. In order to prove that it is semisimple, we must show that its center is trivial. The previous part of the proof shows that

$$(10.4) \quad \mathfrak{g} = \mathfrak{gl} + \mathfrak{w} + \theta\mathfrak{w} ,$$

where \mathfrak{gl} the Lie algebra of $GL(W, C)$ and $\mathfrak{w} \sim W$ that of \mathcal{W} .

Let $Z = X + t_w + \theta t_{w'}$ be a central element. For every $Y \in \mathfrak{gl}$, we must have

$$\begin{aligned} 0 &= [Z, Y] = [X, Y] + [t_w, Y] + \theta[t_{w'}, \theta Y] \\ &= [X, Y] + [t_w, Y] - \theta[t_{w'}, Y^*] . \end{aligned}$$

Since the decomposition (10.4) is respected, each term on the right-hand side must be zero. In particular, $[Y, t_w] = 0$ for every Y . Since $[Y, t_w] = t_{Yw}$, this implies that w is fixed by all elements of $GL(W, C)$ in the connected component of the identity. Therefore $w = 0$. Similarly, $w' = 0$ so that $Z = X$. But then $[X, t_{w''}] = 0$ for every w'' , and the action of \mathfrak{gl} on W is effective. Therefore $X = 0$.

By (10.4), $\mathfrak{p} = \mathfrak{p} \cap \mathfrak{gl} + (\text{id} - \theta)\mathfrak{w}$ and, by Proposition 9.4, $\mathfrak{p} \cap \mathfrak{gl} = \mathfrak{a} + (\text{id} - \theta)\mathfrak{n}$, where \mathfrak{n} is the Lie algebra of N .

The argument preceding Proposition 9.4 shows that, given $a = (a_0, \dots, a_n) \in \mathcal{A}$, $ana^{-1} = (a_i/a_j)n$ when $n \in N$ is as in (9.3) with λ_{ij} being the only non-zero entry. Moreover, $a\tau_w a^{-1} = \tau_{a_i w}$ for $\tau_w \in \mathcal{W}$ with $w \in W_i$.

If h_i the linear functional on \mathfrak{a} projecting onto the i -th component, this shows that \mathfrak{a} acts on $\mathfrak{n} + \mathfrak{w} = \mathfrak{n}_{\mathcal{G}}$ (the Lie algebra of $N_{\mathcal{G}}$) with weights $h_i - h_j$ ($0 \leq j < i \leq n$)

and h_i ($0 \leq i \leq n$). None of these weights being trivial, it follows that \mathfrak{a} is maximal abelian in \mathfrak{p} and the weights just obtained form a system of positive restricted roots of \mathfrak{a} .

Using the isomorphism

$$(10.5) \quad \begin{aligned} \mathfrak{a} = \{(t_0, \dots, t_n)\} &\longrightarrow \{(\tilde{t}_0, \dots, \tilde{t}_{n+1}) : \sum \tilde{t}_i = 0\} \\ (t_0, \dots, t_n) &\longmapsto \left(t_0 - \frac{1}{n+2} \sum t_i, \dots, t_n - \frac{1}{n+2} \sum t_i, -\frac{1}{n+2} \sum t_i \right), \end{aligned}$$

all the roots take the typical form $\tilde{h}_i - \tilde{h}_j$ ($0 \leq i, j \leq n+1, i \neq j$) of the A_{n+1} -system. The fact that this system is irreducible implies that \mathfrak{g} is simple.

The last part of the statement is now obvious. \square

We recall that a smooth map between two Riemannian manifolds is called *conformal* if its differential at any point is a scalar multiple of an orthogonal transformation.

Theorem 10.4. *The action of collineations on projective C -lines is conformal.*

Proof. The elements of U are obviously conformal. Composing with elements of U , we see that it suffices to prove the statement at points in W .

Suppose that g is a translation in \mathcal{W} . Let $w_0 + Cw$ be a tangent C -line at a point w_0 . By (6.3), if φ is the angle of Cw and Cw_0 , $|X|_{w_0+} = c_{|w_0|, \varphi} |X|$ for all tangent vectors $X \in Cw$. Similarly, $|g_* X|_{g(w_0)+} = |X|_{g(w_0)+} = c_{|g(w_0)|, \psi} |X|$, where ψ is the angle of $Cg(w_0)$ and Cw . This implies the statement for $g \in \mathcal{W}$.

If $g \in GL(W, C)$, Corollary 9.3 implies conformality on lines through 0. Composing with translations, it follows that g is conformal on all lines in W . By the decomposition (10.1), this finishes the proof. \square

Since projective C -lines are spheres of the same dimension as C (that we are assuming strictly greater than one), by Liouville's theorem conformal transformations are expressed by fractional linear transformations. For $C = \mathbb{R}$, Theorem 10.4 can be reformulated by replacing the word "conformal" with "fractional linear". This also suggests that, when V is trivial, the group \mathcal{G} that naturally replaces the collineation group is the conformal group when $C \neq \mathbb{R}$, and the group of fractional linear transformations when $C = \mathbb{R}$.

Finally, the collineation group is the group of "basic transformations" of [Te], characterized (when CPW is not a sphere) by the property of mapping Helgason spheres into Helgason spheres.

11. COMPACT RANK-ONE SPACES AND SYMMETRIC CONES

We show that the groups \mathcal{G} are exactly the groups such that $\mathcal{G} \times \mathbb{R}^+$ is the automorphism group of an irreducible symmetric cone [FK]. In particular, \mathcal{G}/U can be imbedded as a domain in real projective space in such a way that \mathcal{G} acts by projective maps.

Taking into account Theorem 10.3, our statement will follow from the following theorem, whose proof is based on representation theory.

Theorem 11.1. *Let \mathfrak{g} be a real simple non-compact Lie algebra whose restricted root system is of type A_ℓ and let G be the adjoint group. Then $\tilde{G} = G \times \mathbb{R}^+$ is the automorphism group of an irreducible symmetric cone.*

The converse is well known, e.g. [FK], p. 108.

Proof. We write $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ for the Cartan decomposition and denote by θ the Cartan involution. We choose $\mathfrak{a} \subset \mathfrak{p}$ maximal abelian, and we complete its complexification $\mathfrak{a}^{\mathbb{C}}$ to a θ -invariant Cartan subalgebra $\mathfrak{h}^{\mathbb{C}}$ of $\mathfrak{g}^{\mathbb{C}}$. We identify $\mathfrak{h}^{\mathbb{C}}$ with its own dual under the Killing form $(\cdot | \cdot)$ and write \mathfrak{h} for the real span of the roots. We also consider $\tilde{\mathfrak{g}} = \mathfrak{g} + \mathbb{R}$, $\tilde{\mathfrak{a}} = \mathfrak{a} + \mathbb{R}$, etc., and extend $(\cdot | \cdot)$ to an inner product on $\tilde{\mathfrak{h}}$ so that \mathbb{R} is orthogonal to \mathfrak{h} . The theory of roots, restricted roots, weights, etc. still applies to the reductive algebra $\tilde{\mathfrak{g}}$. If μ is a weight on $\tilde{\mathfrak{h}}$, we denote by $\bar{\mu}$ its restriction to $\tilde{\mathfrak{a}}$.

We choose an orthonormal basis $\{\varepsilon_1, \dots, \varepsilon_{\ell+1}\}$ of $\tilde{\mathfrak{a}} \subset \tilde{\mathfrak{h}}$ so that the restricted roots of $\tilde{\mathfrak{g}}$ are $\bar{\varepsilon}_i - \bar{\varepsilon}_j$ ($i \neq j$). Since the Weyl group is transitive on the roots, the corresponding root spaces $\mathfrak{g}^{\bar{\varepsilon}_i - \bar{\varepsilon}_j}$ have the same dimension, to be denoted by d . The \mathbb{R} -part of $\tilde{\mathfrak{a}}$ is spanned by $\sum \varepsilon_i$. We fix $\bar{\varepsilon}_i - \bar{\varepsilon}_j$ with $i < j$ as the positive roots.

Let (ρ, V) be the irreducible representation of $\tilde{\mathfrak{g}}$ with highest weight $2\varepsilon_1$. By Theorem 4.12 in [Hel3], such a representation exists and it is a spherical representation, i.e. V has a K -fixed vector. Indeed, $2\varepsilon_1$ is in $\tilde{\mathfrak{a}}$ and it satisfies the integrality condition

$$\frac{(2\varepsilon_1 | \varepsilon_i - \varepsilon_j)}{(\varepsilon_i - \varepsilon_j | \varepsilon_i - \varepsilon_j)} \in \mathbb{Z} \quad (\forall i \neq j).$$

We claim that $\dim_{\mathbb{C}} V = \dim_{\mathbb{R}} \tilde{G}/K$. To see this, we compute the dimensions of the \mathfrak{a} -weight spaces $\bar{V}_{\bar{\mu}} = \{v \in V : \rho(H)v = \bar{\mu}(H)v, \forall H \in \mathfrak{a}\}$. Each $\bar{V}_{\bar{\mu}}$ is the sum of the \mathfrak{h} -weight spaces $V_{\mu'} = \{v \in V : \rho(H)v = \mu'(H)v, \forall H \in \mathfrak{h}\}$, for those \mathfrak{h} -weights μ' whose restriction to \mathfrak{a} is $\bar{\mu}$.

By the general theory (cf. [Hum], p. 108, 114), the \mathfrak{h} -weights μ of ρ have the form $2\varepsilon_1 - \alpha$, where α is a positive \mathfrak{h} -root and $(\alpha | 2\varepsilon_1) > 0$. It follows that the only \mathfrak{h} -weight of ρ which restricts to $2\bar{\varepsilon}_1$ is $2\varepsilon_1$. Hence $\bar{V}_{2\bar{\varepsilon}_1} = V_{2\varepsilon_1}$ is one-dimensional.

By action of the Weyl group, $\bar{V}_{2\bar{\varepsilon}_j}$ is one-dimensional for $1 \leq j \leq \ell + 1$. The other \mathfrak{a} -weights are contained in the convex hull of the $2\bar{\varepsilon}_j$ and equal to $2\bar{\varepsilon}_j$ minus a sum of simple restricted roots. Therefore the other \mathfrak{a} -weights can only be $\bar{\varepsilon}_i + \bar{\varepsilon}_j$, with $i < j$. By invariance under the Weyl group again, they must have the same multiplicity, so that we can restrict our attention to the weight $\bar{\varepsilon}_1 + \bar{\varepsilon}_2$.

It is clear that $\bar{V}_{\bar{\varepsilon}_1 + \bar{\varepsilon}_2}$ is the direct sum of all $V_{2\varepsilon_1 - \alpha}$ with $\bar{\alpha} = \bar{\varepsilon}_1 - \bar{\varepsilon}_2$. We will prove that each such $V_{2\varepsilon_1 - \alpha}$ equals $\rho(\mathfrak{g}_{-\alpha})V_{2\varepsilon_1}$, with $\mathfrak{g}_{-\alpha}$ the \mathfrak{h} -root space relative to $-\alpha$, hence is one-dimensional. This will then show that $\dim \bar{V}_{\bar{\varepsilon}_1 + \bar{\varepsilon}_2} = d$.

By [Hum], p.108, $V_{2\varepsilon_1 - \alpha}$ is spanned by the vectors

$$(11.1) \quad v = \rho(X_{-\alpha_k}) \cdots \rho(X_{-\alpha_1})v^+,$$

where v^+ is a fixed non-zero element in $V_{2\varepsilon_1}$, the α_j are positive \mathfrak{h} -roots, $\sum_{i=1}^k \alpha_i = \alpha$ and $X_{-\alpha_i}$ spans the \mathfrak{h} -root space $\mathfrak{g}_{-\alpha_i}$. We may assume that $v \neq 0$.

Since each $\bar{\alpha}_i$ is a positive restricted root or zero, there is one i_0 for which $\bar{\alpha}_{i_0} = \bar{\varepsilon}_1 - \bar{\varepsilon}_2$, while $\bar{\alpha}_i = 0$ for $i \neq i_0$. If $\bar{\alpha}_1 = 0$, then $2\varepsilon_1 - \alpha_1$ is not a weight for ρ , hence $\rho(X_{-\alpha_1})v^+ = 0$, a contradiction. So $i_0 = 1$. Now we prove that, if $k > 1$, the number of factors in (11.1) can be reduced by one. Indeed, since $\rho(X_{-\alpha_2})v^+ = 0$,

$$\rho(X_{-\alpha_2})\rho(X_{-\alpha_1})v^+ = \rho([X_{-\alpha_2}, X_{-\alpha_1}])v^+ = \rho(X_{-\alpha'_1})v^+,$$

with some $X_{-\alpha'_1} \in \mathfrak{g}_{-\alpha'_1}$, $\alpha'_1 = \alpha_1 + \alpha_2$.

Repeating this argument we find that v is in $\rho(\mathfrak{g}_{-\alpha})V_{2\varepsilon_1}$.

By invariance under the Weyl group, $\dim_{\mathbb{C}} \bar{V}_{\bar{\varepsilon}_i + \bar{\varepsilon}_j} = d$ for all $i < j$, and adding up the dimensions, we find that $\dim_{\mathbb{C}} V = \ell + 1 + \frac{\ell(\ell+1)}{2}d$. In $\tilde{\mathfrak{g}} = \mathfrak{k} + \tilde{\mathfrak{p}}$ we have $\tilde{\mathfrak{p}} = \tilde{\mathfrak{a}} + \sum_{i < j} (\text{id} - \theta)\mathfrak{g}_{\bar{\varepsilon}_i - \bar{\varepsilon}_j}$. Counting the dimensions, we find that $\dim_{\mathbb{C}} V = \dim_{\mathbb{R}} \tilde{G}/K$.

We imbed \tilde{G}/K into V by choosing a K -invariant vector e and defining the map $gK \mapsto \rho(g)e$. Then $V^0 = \rho(\tilde{\mathfrak{p}})e$ is the tangent space of $\Omega = \rho(\tilde{G})e$ at e (under the usual identification). V^0 is a real form of V and it is $\rho(K)$ -invariant. Introducing an inner product invariant under the compact form of $G^{\mathbb{C}}$, $\rho(\tilde{\mathfrak{p}})$ consists of Hermitian linear transformations. Then $\rho(\tilde{\mathfrak{a}}), \rho(\exp \tilde{\mathfrak{a}})$ are simultaneously diagonalizable and real. It follows that $\Omega = \rho(K)\rho(\tilde{A})e$ is in V^0 . Since its tangent space at e is all of V^0 , it is open. Since the \mathbb{R}^+ -part of \tilde{G} acts by positive scalar transformations, Ω is a cone in V^0 .

To see that Ω is symmetric in the sense of [FK], i.e. self-dual, we observe that our choice of inner product guarantees that $\rho(\tilde{G})$ is closed under taking adjoints. This is enough to prove that Ω is self-dual, cf. [FK], p. 20, Exercise 8. \square

One can also describe Ω in more detail. By the Cartan decomposition, $V^0 = \rho(K)V^{\text{rad}}$, where V^{rad} is the subspace $\rho(\tilde{\mathfrak{a}})e$. Properly normalizing the inner product, the vectors $e_i = \rho(\varepsilon_i)e$ form an orthonormal basis of V^{rad} . Then $\Omega^{\text{rad}} = \Omega \cap V^{\text{rad}}$ is just the positive quadrant in V^{rad} , and $\Omega = \rho(K)\Omega^{\text{rad}}$.

We also note that the symmetric space G/K can be realized as $\rho(K)\rho(\exp_G \mathfrak{a})e$, where $\rho(\exp_G \mathfrak{a})e$ is the hyperboloid $\{\sum_1^{\ell+1} t_i e_i : t_1 t_2 \cdots t_{\ell+1} = 1\}$ in V^{rad} .

Being a dual cone, Ω is convex. Its extremal generators can only be the $\rho(K)$ -images of the edges in Ω^{rad} and K acts transitively on these. So \mathcal{E}_1 , the intersection of the set of extremal generators with the unit sphere, is just $K \cdot e_1$. The stabilizer of e_1 in K is the same as the centralizer of ε_1 in K , and this is the M' -part of the Langlands decomposition of the parabolic subgroup $P = M'A'N'$ determined by the simple roots $\varepsilon_i - \varepsilon_{i+1}$ with $2 \leq i \leq \ell$.

We identify e_1 and \mathcal{E}_1 with their images $\tilde{e}_1, \tilde{\mathcal{E}}_1$ in the projective space PV^0 and write $\tilde{\rho}$ for the action of G on PV^0 induced by ρ . We claim that $\tilde{\rho}(G)\tilde{e}_1 = \tilde{\mathcal{E}}_1$ and the stabilizer of \tilde{e}_1 is P . For this, we consider $\rho(\exp t\varepsilon_1)e = e^t e_1 + e_2 + \cdots + e_{\ell+1}$, which is fixed under the conjugate $K^{\exp t\varepsilon_1}$ of K . Passing to PV^0 and letting $t \rightarrow \infty$, \tilde{e}_1 is fixed under the limit of $K^{\exp t\varepsilon_1}$, which is $M'N'$, and also under A' , which acts on e_1 by scalars. This shows that $\tilde{\mathcal{E}}_1$ is one of the Satake-Fürstenberg boundaries of G/K .

It is a result of U. Hirzebruch (cf. [FK], p.78, Exercise 5) that, for any irreducible symmetric cone, \mathcal{E}_1 is a compact symmetric space of rank one, and (by classification) every such space arises in this way. We can now reprove this result, and a little more, if we show that, starting with any compact rank-one symmetric space, realizing it as in Section 5, then using Theorem 11.1 to construct the corresponding cone and its \mathcal{E}_1 , we get back to the initial space.

For this, it is enough to check that when we apply Theorem 11.1 with \mathcal{G} and U in place of G and K , the group M' will be the same as the group K of Sections 4-10. Now, in the Lie algebra of it, we can take \tilde{h}_{n+1} as the element corresponding to ε_1 . After the coordinate change (10.5), this is a scalar multiple of $h_0 + \cdots + h_n$, which generates the group of scalar transformations of W . The stabilizer of this in

U is indeed K , by Corollary 6.5.

APPENDIX. THE NON-COMPACT SYMMETRIC SPACES

What follows is redoing [CDKR2] from a different starting point, to make it more compatible with the approach to compact spaces that we have taken.

Let B_W be the open unit ball in W . The tangent space $T_w B_W$ at a point in $w \in B_W$ is naturally identified with W itself. We introduce a Riemannian metric on B_W by assigning, on the tangent space $T_w B_W$ at $w \in B_W$, the scalar product $\langle \cdot, \cdot \rangle_{w-}$ such that

$$(A.1) \quad \langle X, Y \rangle_{w-} = \begin{cases} \frac{\langle X, Y \rangle}{(1 - |w|^2)^2} & \text{if } X, Y \in Cw, \\ \frac{\langle X, Y \rangle}{1 - |w|^2} & \text{if } X, Y \in (Cw)^\perp, \\ 0 & \text{if } X \in Cw, Y \in (Cw)^\perp, \end{cases}$$

for $w \neq 0$ and, passing to the limit for $w \rightarrow 0$,

$$(A.2) \quad \langle X, Y \rangle_{0-} = \langle X, Y \rangle \quad \text{for every } X, Y.$$

In [CDKR2] these formulas are obtained towards the end, in Section 7; here they are definitions. Note that in [CDKR2] this metric is multiplied by a factor 4. This multiplies by a constant the arc lengths and curvatures, but does not change anything essential such as isometries or geodesic submanifolds.

It is immediate that the same K of Section 4 consists of isometries fixing the origin. Adapting the proofs in Section 6, one easily obtains the following properties.

Lemma A.1. *The geodesic γ going through the origin with tangent vector $w \in S_W$ is the diameter in the direction of w , parametrized as $\gamma(t) = (\tanh t)w$. The group of isometries of B_W fixing the origin is K .*

The maps a_t defined by

$$(A.3) \quad a_t(\zeta, v) = ((\sinh t \zeta + \cosh t \mathbf{1})^{-1} (\cosh t \zeta + \sinh t \mathbf{1}), (\sinh t \zeta + \cosh t \mathbf{1})^{-1} v).$$

form a one-parameter group A of isometries of B_W . This can be proved by a computation very similar to the proof of Lemma 6.2.

Since the orbit of $(0, 0)$ under A consists of the points $a_t(0, 0) = (\tanh t, 0)$, it follows that the group G of isometries of B_W is transitive on B_W . We also see that $G = KAK$. Since the isometry $w \mapsto -w$ has a single fixed point, it also follows that B_W is a symmetric space. Transitivity of K on S_W implies that the rank of B_W is one.

Simple modifications to the proof of Theorem 7.2 give that the sectional curvature of a plane element E spanned by $X, Y \in W$ in the tangent space to B_W at the origin is the negative of $\sigma(E)$ in (7.1).

The unit balls B_1 in $W_1 = C_1 \oplus V_1$ and B_2 in $W_2 = C_2 \oplus V_2$ are isometric if and only if $(C_1, V_1) \sim (C_2, V_2)$ as C -modules. However, the degenerate cases (\mathbb{R}, V) and $(\mathbb{R} \oplus V, 0)$ give different models of the same space. The map

$$(t + v, 0) \mapsto \left(\frac{2t}{1 + t^2 + |v|^2}, \frac{2v}{1 + t^2 + |v|^2} \right)$$

from the second to the first is an isometry up to a factor 2. For $V \neq 0$, these are respectively the Klein model and the Poincaré model of real hyperbolic space. For $V = 0$, they trivially become models of the real line. [CDKR2] contains a classification-independent proof of the fact that the metrics (A.1) give all the rank-one symmetric spaces of the non-compact type (together with \mathbb{R} , the irreducible Euclidean non-compact symmetric space).

The maps a_t in (A.3) extend to homeomorphisms of CPW . The extension to $W \setminus \{(\zeta, v) : \zeta \neq -\coth t\}$ is obvious, and the extension to the rest of CPW is, in analogy with (6.10),

$$\begin{aligned} a_t(-\coth t, v) &= [\mathbf{1}, -(\sinh t)v] , \\ a_t([\mathbf{1}, v]) &= (\coth t, (\sinh t)^{-1}v) \\ a_t([0, v]) &= [0, v] . \end{aligned}$$

Therefore, all isometries of B_W extend to homeomorphisms of CPW . In analogy with Corollary 7.4, it follows that these extensions are collineations.

To tie up with [CDKR2], we use the Cayley transform c (cf. [CDKR2], p. 208), which in our notation is

$$(\zeta', v') = c(\zeta, v) = ((\mathbf{1} - \zeta)^{-1}(\mathbf{1} + \zeta), 2(\mathbf{1} - \zeta)^{-1}v) .$$

Clearly,

$$\Re \zeta' - \frac{1}{4}|v'|^2 = \frac{1 - |\zeta|^2 - |v|^2}{|\mathbf{1} - \zeta|^2} ,$$

which shows that c maps B_W onto

$$D = \{(\zeta', v') : \Re \zeta' - \frac{1}{4}|v'|^2 > 0\} .$$

Define $B : V \times V \rightarrow C$ by

$$\langle B(v, v'), \zeta \rangle = \langle \zeta v, v' \rangle .$$

Among the isometries of D , with the metric induced on it by c , one has the “translation” group $\tilde{N} = \{\tilde{n}_{(z,u)} : z \in C', u \in V\}$, with

$$\tilde{n}_{(z,u)}(\zeta, v) = \left(\zeta + z + \frac{1}{2}B(v, u) + \frac{1}{4}|u|^2, v + u \right) ,$$

acting simply transitively on the level sets of the height function $h(\zeta, v) = \Re \zeta - \frac{1}{4}|v|^2$.

The Lie algebra \mathfrak{n} of \tilde{N} is an H -type Lie algebra satisfying the J^2 -condition. The discussion in Section 3 shows that all such algebras show up in this way.

If $\tilde{a}_t = ca_t c^{-1}$, then

$$\tilde{a}_t(\zeta, v) = (e^{2t}\zeta, e^t v) ,$$

the group $\tilde{A} = cAc^{-1}$ normalizes \tilde{N} , and $\tilde{N}\tilde{A}$ acts simply transitively on D . This is the starting point in [CDKR2].

We note that c is almost the same as $b_{\frac{\pi}{4}}$ defined in Lemma 6.2, except that the factor 2 in c is replaced by $\sqrt{2}$ in $b_{\frac{\pi}{4}}$. One can in fact replace c by $b_{\frac{\pi}{4}}$ and D by $D' = \{(\zeta', v') : \Re \zeta' - \frac{1}{2}|v'|^2 > 0\}$, a choice that may be preferable in some ways.

REFERENCES

- [Ba] Baez, J., *The octonions*, Bull. Amer. Math. Soc. **39** (2002), 145-205.
- [Be] Besse, A., *Manifolds all of whose geodesics are closed*, Springer Verlag, 1978.
- [CDKR1] Cowling, M., Dooley, A. H., Korányi, A., Ricci, F., *H-type groups and Iwasawa decompositions*, Adv. Math. **87** (1991), 1-41.
- [CDKR2] Cowling, M., Dooley, A. H., Korányi, A., Ricci, F., *An approach to symmetric spaces of rank one via groups of Heisenberg type*, J. Geom. Anal. **8** (1998), 199-237.
- [FK] Faraut, J., Korányi, A., *Analysis on symmetric cones*, Oxford Univ. Press, 1994.
- [Hei] Heintze, E., *On homogeneous manifolds of negative curvature*, Math. Ann. **21** (1974), 23-34.
- [Hel1] Helgason, S., *Differential geometry, Lie groups and symmetric spaces*, Acad. Press, 1978.
- [Hel2] Helgason, S., *Totally geodesic spheres in compact symmetric spaces*, Math. Ann. **165** (1966), 309-317.
- [Hel3] Helgason, S., *Geometric analysis on symmetric spaces*, Amer. Math. Soc., Math. Surveys 39, 1994.
- [Hum] Humphreys, J. E., *Introduction to Lie algebras and representation theory*, Springer Verlag, 1972.
- [Hus] Husemoller, D., *Fiber bundles*, Springer Verlag, 1994.
- [K] Kaplan, A., *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc. **259** (1980), 147-153.
- [KN] Kobayashi, S., Nomizu, K., *Foundations of differential geometry, I and II*, Wiley Intersc., 1996.
- [KR] Korányi, A., Ricci, F., *A classification-free construction of rank-one symmetric spaces*, Bull. Kerala Math. Assoc., Special Issue (2005), 72-88.
- [M] Mostow, G. D., *Strong rigidity of locally symmetric spaces*, Princeton Univ. Press, 1973.
- [R] Riehm, C., *The automorphism group of a composition of quadratic forms*, Trans. Amer. Math. Soc. **269** (1982), 403-414.
- [S] Springer, T. A., *The projective octave plane. I, II*, Indag. Math. **22** (1960), 74-101.
- [Ta] Takahashi, R., *Quelques résultats sur l'analyse harmonique dans l'espace symétrique non compact de rang 1 du type exceptionnel*, Lecture Notes in Math, Springer, 739 (1979), 511-567.
- [Te] Takeuchi, M., *Basic transformations of symmetric R-spaces*, Osaka J. Math. **25** (1988), 259-297.
- [W] Wolf, J., *Spaces of constant curvature*, McGraw-Hill, 1967.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, LEHMAN COLLEGE, CITY UNIVERSITY OF NEW YORK, BRONX, NY 10468-1589, USA

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, 56126 PISA, ITALY